DOI: 10.1002/aimg.b.32962

RESEARCH ARTICLE

Latin American Trans-ancestry INitiative for OCD genomics (LATINO): Study protocol

James J. Crowley^{1,2} | Carolina Cappi^{3,4} | Marcos E. Ochoa-Panaifo⁵ | Renee M. Frederick⁶ | Minjee Kook⁶ | Andrew D. Wiese⁶ | Diana Rancourt⁷ Elizabeth G. Atkinson⁸ | Paola Giusti-Rodriguez⁹ | Jacey L. Anderberg⁶ | Latin American Trans-ancestry INitiative for OCD genomics, Brazilian Obsessive-Compulsive Spectrum Disorder Working Group | Jonathan S. Abramowitz¹⁰ | Victor R. Adorno¹¹ | Cinthia Aguirre¹² | Gilberto S. Alves¹³ | Gustavo S. Alves^{14,15} | NaEshia Ancalade¹ | Alejandro A. Arellano Espinosa¹⁶ | Paul D. Arnold¹⁷ | Daphne M. Ayton⁶ | Izabela G. Barbosa¹⁸ | Laura Marcela Barón Castano¹⁹ | Cynthia N. Barrera²⁰ | María Celeste Berardo²¹ | Dayan Berrones²² | John R. Best²³ | Tim B. Bigdeli^{24,25} | Christie L. Burton²⁶ | Joseph D. Buxbaum³ | Jennifer L. Callahan²⁷ | Maria Cecília B. Carneiro²⁸ | Sandra L. Cepeda²⁹ | Evelyn Chazelle²¹ | Jessica M. Chire³⁰ | Macarena Churruca Munoz³¹ | Pamela Claisse Quiroz³² | Journa Cobite³³ | Jonathan S. Comer³⁴ | Daniel L. Costa⁴ | Jennifer Crosbie^{26,35} | Victor O. Cruz^{36,37} | Guillermo Dager³⁸ | Luisa F. Daza¹⁹ | Anabel de la Rosa-Gómez³⁹ | Daniela del Río⁴⁰ | Fernanda Z. Delage⁴¹ Carolina B. Dreher^{42,43} | Lucila Fay²¹ | Tomas Fazio²¹ | Ygor A. Ferrão⁴⁴ | Gabriela M. Ferreira^{41,45} | Edith G. Figueroa⁴⁶ | Leonardo F. Fontenelle^{47,48} | Diego A. Forero⁴⁹ | Daniele T. H. Fragoso⁴¹ | Bharathi S. Gadad⁵⁰ | Sheldon R. Garrison⁵¹ I Andres González⁵² | Laura D. Gonzalez²¹ | Marco A. González³⁹ | Polaris Gonzalez-Barrios^{53,54} | Wayne K. Goodman⁶ | Dorothy E. Grice³ | Jerry Guintivano² | Daniel G. Guttfreund⁵⁵ | Andrew G. Guzick⁶ | Matthew W. Halvorsen¹ | Joseph D. Hovey⁵⁶ | Hailiang Huang⁵⁷ | Jonathan Irreño-Sotomonte⁵⁸ | Reinhard Janssen-Aguilar⁵⁹ | Matias Jensen⁶⁰ | Alexandra Z. Jimenez Reynolds⁶ | Joali Alexandra Juárez Lujambio⁶¹ | Nasim Khalfe⁶² | Madison A. Knutsen^{6,63} | Caleb Lack⁶⁴ | Nuria Lanzagorta⁶⁵ | Monicke O. Lima⁴ | Melanie O. Longhurst⁵⁰ |

The views expressed in this article do not necessarily represent the views of the National Institutes of Health, the Department of Health and Human Services, or the United States Government. For affiliations refer to page 11

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics published by Wiley Periodicals LLC.

David A. Lozada Martinez⁵ | Elba S. Luna³⁶ | Andrea H. Margues⁶⁶ | Molly S. Martinez^{67,68} | Maria de Los Angeles Matos²¹ | Caitlyn E. Maye⁶ | Joseph F. McGuire⁶⁹ | Gabriela Menezes⁷⁰ | Charlene Minava⁷¹ | Tomás Miño⁶⁰ | Sara M. Mithani⁷² | Circe Montes de Oca⁷³ | Alonso Morales-Rivero⁷⁴ Maria E. Moreira-de-Oliveira⁷⁰ Olivia J. Morris⁶ Sandra I. Muñoz³⁹ | Zainab Naggash²³ | Ambar A. Núñez Bracho⁵² | Belinda E. Núñez Bracho⁵² | Maria Corina Ochoa Rojas¹⁹ | Luis A. Olavarria Castaman⁵ | Trinidad Olivos Balmaceda⁷⁵ | Iliana Ortega¹⁷ | Darpan I. Patel⁷² | Ainsley K. Patrick⁶⁹ | Mariel Paz y Mino^{76,77} | Jose L. Perales Orellana⁷⁸ | Bárbara Perdigão Stumpf¹⁸ | Tamara Peregrina⁷³ | Tania Pérez Duarte²⁰ | Kelly L. Piacsek⁵¹ | Maritza Placencia⁷⁹ | María Belén Prieto²¹ | Lucas C. Quarantini^{14,80} | Yana Quarantini-Alvim^{14,81} | Renato T. Ramos^{35,82} | Iaroslava C. Ramos^{82,83} | Vanessa R. Ramos⁴ | Keslev A. Ramsev⁶⁹ | Elise V. Ray¹ | Margaret A. Richter^{82,84} | Bradley C. Riemann⁵¹ | Juan C. Rivas^{19,85,86,87} | Maria C. Rosario⁸⁸ | Camilo J. Ruggero²⁷ | Angel A. Ruiz-Chow⁷⁴ | Alejandra Ruiz-Velasco⁵⁰ | Melisa N. Sagarnaga⁸⁹ | Aline S. Sampaio^{14,15,80} | Leonardo C. Saraiva⁴ | Russell J. Schachar^{26,35} | Sophie C. Schneider⁶ | Ethan J. Schweissing⁶ | Laura D. Seligman⁵⁶ | Roseli G. Shavitt⁴ | Keaton J. Soileau⁶ | S. Evelyn Stewart^{23,90,91} | Shaina B. Storch⁹² | Emily R. Strouphauer⁶² | Vissente Tapia Cuevas⁷⁵ | Kiara R. Timpano²⁹ | Beatriz Treviño-de la Garza⁹³ | Alexie Vallejo-Silva⁵⁸ | Javier Vargas-Medrano⁵⁰ | María I. Vásquez⁹⁴ | Guadalupe Vidal Martinez⁵⁰ | Saira A. Weinzimmer⁶ | Mauricio A. Yanez⁶ | Gwyneth Zai^{84,95} | Lina M. Zapata-Restrepo^{87,96,97} | Luz M. Zappa^{21,98,99} Raquel M. Zepeda-Burgos¹⁰⁰ | Anthony W. Zoghbi^{6,101} | Euripedes C. Miguel⁴ Carolyn I. Rodriguez^{102,103} | Mayra C. Martinez Mallen⁷⁴ | Pablo R. Moya^{31,104} Tania Borda^{105,106} | María Beatriz Moyano^{21,107,108} | Manuel Mattheisen^{109,110} Stacev Pereira¹¹¹ Gabriel Lázaro-Muñoz^{112,113} Karen G. Martinez-Gonzalez⁵³ Michele T. Pato¹¹⁴ | Humberto Nicolini^{115,116} | Eric A. Storch⁶

Correspondence

James J. Crowley, Departments of Genetics and Psychiatry, University of North Carolina at Chapel Hill, 120 Mason Farm Road, Chapel Hill, NC, 27599, USA. Email: crowley@unc.edu

Eric A. Storch, Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, 1977 Butler Blvd, Suite 4-400, Houston, TX, 77030, USA. Email: eric.storch@bcm.edu

Abstract

Obsessive-compulsive disorder (OCD) is a debilitating psychiatric disorder. Worldwide, its prevalence is $\sim 2\%$ and its etiology is mostly unknown. Identifying biological factors contributing to OCD will elucidate underlying mechanisms and might contribute to improved treatment outcomes. Genomic studies of OCD are beginning to reveal long-sought risk loci, but >95% of the cases currently in analysis are of homogenous European ancestry. If not addressed, this Eurocentric bias will result in OCD genomic findings being more accurate for individuals of European ancestry than

Funding information

National Institute of Mental Health, USA. Grant/Award Numbers: K23MH121669. U01MH125062, U01MH125050; Caroline Wiess Law Fund for Research in Molecular Medicine, Grant/Award Number: K01 MH121659; Conselho Nacional de Desenvolvimento Científico e. Tecnológico, Brazil, Grant/Award Number: 302526/2018-8: Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro, Brazil, Grant/Award Number: CNE E-26/203.052/2017; Hispanic Clinical and Translational Research Education and Career Development (HCTRECD), Grant/Award Number: R25MD007607; National Institute of Developmental Psychiatry for Children and Adolescent, Grant/Award Number: Fapesp 2014/50917-0.CNPq 465550/2014-2; National Center for Complementary and Integrative Health, Grant/Award Number: 1R34AT010661-0: Beatrice and Samuel A Seaver Foundation the Mindworks Charitable Lead Trust, NIMH, Grant/Award Number: R01MH124679

other ancestries, thereby contributing to health disparities in potential future applications of genomics. In this study protocol paper, we describe the Latin American Trans-ancestry INitiative for OCD genomics (LATINO, https://www.latinostudy.org). LATINO is a new network of investigators from across Latin America, the United States, and Canada who have begun to collect DNA and clinical data from 5000 richly phenotyped OCD cases of Latin American ancestry in a culturally sensitive and ethical manner. In this project, we will utilize trans-ancestry genomic analyses to accelerate the identification of OCD risk loci, fine-map putative causal variants, and improve the performance of polygenic risk scores in diverse populations. We will also capitalize on rich clinical data to examine the genetics of treatment response, biologically plausible OCD subtypes, and symptom dimensions. Additionally, LATINO will help elucidate the diversity of the clinical presentations of OCD across cultures through various trainings developed and offered in collaboration with Latin American investigators. We believe this study will advance the important goal

KEYWORDS

genetic, genomic, GWAS, Latin American, obsessive-compulsive disorder, OCD

of global mental health discovery and equity.

1 | RATIONALE

1.1 | Overview

Genomics is now identifying long-elusive risk genes, distinguishing related disorders, and beginning to guide therapy. Unfortunately, most of these advances primarily benefit individuals of European ancestry (EA), potentially exacerbating existing health disparities (Martin et al., 2019). Furthermore, admixed populations, such as Latino/Latina (hereafter Latino) populations, constitute more than a third of the U.S. population and yet are markedly understudied in medical and psychiatric genetics (Peterson et al., 2019). The reasons for the current Eurocentric bias in genomics include sociocultural factors (Sanderson et al., 2013), such as more funding opportunities in higher-income countries, and analytical reasons where non-EA samples are often simply excluded from EA-focused analyses. Correcting this disparity will expand our understanding of disease risk from common variation observed among individuals of all ancestries, with particular benefit to understudied Latino individuals. For example, novel loci are bound to be discovered owing to alleles specific to Latino populations, which will expand our view of disease pathophysiology. Existing loci can be fine mapped by taking advantage of linkage disequilibrium differences across ancestries, bringing us closer to identifying causal variants. In addition, polygenic risk scores (PRS: a way of measuring common genetic risk factors) will be more accurate in non-EA individuals if the discovery sample includes their ancestral background. Overall, greater diversity needs to be sampled to ensure that personalized medicine and pharmacogenetics will not lead to even greater health disparities in genomics.

Obsessive-compulsive disorder (OCD) is a disorder that needs to both markedly diversify its genomic sample and to increase its sample size overall. In the past year, the OCD genome-wide association study (GWAS) sample size has grown from just 2688 cases to \sim 50,000 cases (\sim 10,000 from clinics, \sim 10,000 from biobanks, and \sim 30,000 self-reported OCD cases from 23andMe), according to the latest, unpublished freeze from the Psychiatric Genomics Consortium (PGC) OCD Working Group (presented in October 2022 at the World Congress of Psychiatric Genetics in Florence, Italy). Unfortunately, the sample is far from diverse - over 95% of the cases are of EA. Preliminary GWAS results, based on EA cases and controls, have yielded the first 30 genome-wide significant hits for OCD, and brought OCD GWAS to the point where many more loci will be discovered as sample size increases. One of the next steps will be to perform trans-ancestry meta-analyses; but with just \sim 2000 non-EA cases from all over the world, power for these analyses is low. Therefore, we will collect a large number of non-EA participants and work together with PGC OCD to perform the most highly powered analysis possible. Our goal is to reduce health disparities and identify both common and ancestry-specific loci for OCD.

Including people of Latino ancestry from areas in North and South America provides a fertile ground for such a collection. The genetic makeup of Latino individuals has been shaped by a history of extensive admixture between Africans, Europeans, and indigenous peoples (Norris et al., 2018). Thus, the region is ideal to examine the interplay of genetic factors from these three major ancestral groups. Therefore, we assembled a network of over 50 sites in Latin America, the United States, and Canada that treat a large number of ancestrally diverse OCD patients (see Figure 1). We have come together with experts in the clinical aspects of OCD, OCD genetics and the genetics





FIGURE 1 LATINO study sites.

of admixed populations for a new NIH-funded study called the Latin American Trans-ancestry INitiative for OCD genomics (LATINO, https://www.latinostudy.org).

First, LATINO will collect DNA and comprehensive clinical data from at least 5000 Latino OCD cases. These cases will be matched with controls collected by other Latino psychiatric genetics projects. Second, the combined cohort of cases and controls will be subject to genome-wide single nucleotide polymorphism (SNP) genotyping, allowing us to perform a novel trans-ancestry GWAS in collaboration with the PGC OCD Working Group. Third, we will leverage these diverse study data to fine-map new OCD risk loci and compare the performance of the newly diverse PRS with EA-only PRS. We believe this study will highlight the importance of including all major ancestral groups in genomic studies, both as a strategy to improve power to find disease associations, as well as to reduce health disparities, in potential future applications of genomics in precision medicine.

LATINO, however, is much more than a genetic study. OCD is under-recognized, under-diagnosed, under-treated, and under-studied in Latino individuals, contributing to health disparities (Abramowitz et al., 2017; Benuto, 2017; Perez et al., 2022; Wetterneck et al., 2012). In addition to decreasing the Eurocentric bias in OCD genetics, LATINO is working with stakeholders and patient-centered organizations to increase awareness of OCD in Latino individuals and improve their access to evidence-based treatments. For example, LATINO is playing a central role in executing social media-based awareness campaigns and providing hands-on training for psychologists and psychiatrists interested in treating OCD. LATINO is also facilitating OCD research in Latin America by translating key instruments into Spanish and Portuguese, validating them in Latino individuals, and providing a genomics training program. Finally, LATINO is helping Latin American OCD researchers identify funding opportunities and establish mutually beneficial collaborations. To that end, LATINO will actively contribute to decreasing disparities on multiple levels.

1.2 | Background

OCD is a neuropsychiatric disorder characterized by recurrent, unwanted thoughts, ideas, images or urges and/or repetitive behaviors that are frequently performed in response to the distress associated with the obsessions (American Psychiatric Association, 2013; World Health Organization, 1992). Individuals with OCD experience exaggerated concerns about danger, hygiene or harm that result in persistent conscious attention to the perceived threat (obsessions) (Goodman et al., 2021). In response to distress associated with obsessions, the individual performs compulsions to neutralize the anxiety/distress, which provides temporary relief. However, relief is negatively reinforcing, leading to repetitive, compulsive behavior when obsessions inevitably recur (Pauls et al., 2014). OCD is a multidimensional disorder shown to consist of roughly four primary symptom dimensions (contamination, symmetry, forbidden thoughts and related compulsions, checking) (Bloch et al., 2008) which may have distinct neural circuitry (Mataix-Cols et al., 2004), genetic (Hasler et al., 2007), and etiological origins (Leckman et al., 2009). Additionally, some subtypes such as early-onset and tic-related OCD may be etiologically meaningful subtypes of the disorder (Brander et al., 2021; Leckman et al., 2010).

The OCD lifetime prevalence is ~2% worldwide (Karno et al., 1988; Ruscio et al., 2010; Wetterneck et al., 2012), with onset often in childhood and similar prevalence by sex. Studies of OCD in Latin America have shown similar prevalences (Wetterneck et al., 2012). Most OCD cases have a comorbid psychiatric disorder (e.g., tic, mood, and anxiety disorders) (Fullana et al., 2009). Patients with OCD are at substantial risk of suicide (~10 times higher than the population prevalence) (Fernández de la Cruz et al., 2017). OCD was ranked among the 10 leading causes of years lost to disability in 2008 (Mathers & World Health Organization, 2008). Although medication and behavioral therapy are useful (McGuire et al., 2015), symptom control is imperfect, and the course is often chronic.

TABLE 1 LATINO study specific aims.

Aim 1. Diversify OCD genomics sample

A. Collect saliva-derived DNA from 5000 Latino OCD cases from a network of clinics across Latin America, the United States and Canada. Matched controls are available through collaboration.

B Neuropsychiatric Genetics WILEY 5 of 18

B. Perform a detailed phenotypic assessment of each case (e.g., comorbidities, symptom dimensions, severity, and treatment response).

Aim 2. Discover genomic loci for OCD

- A. Genotype all 5000 Latino OCD cases with the Illumina Global Screening Array (GSA) and match with data from at least 10,000 Latino controls.
- B. Perform a GWAS on the 5000 OCD cases and at least 10,000 matched controls and meta-analyze with results from the PGC OCD Working Group.

Aim 3. Fine-map OCD loci and test accuracy of PRS

- A. Evaluate fine-mapping gains in terms of reductions in the number of credible SNPs at genome-wide significant loci following transancestry GWAS meta-analysis.
- B. Compare the performance of PRS trained on our diverse sample with analogous scores based on EA-only samples.

A deeper understanding of the pathophysiology of OCD is critical to the development of more effective treatments. Successful completion of our aims will make important contributions to understanding the etiology of this disorder, alongside diversifying the sample to include understudied populations. We believe that the identification of specific genes and biological pathways mediating susceptibility to OCD will lead to improved detection, differential diagnosis, and treatment strategies (Grassi et al., 2020). Our sampling scheme is not only cost-effective; but also captures representative, comprehensively phenotyped Latino OCD cases in multiple Latin American nations. The participation of trained bioethicists will help guide the collection of samples in a culturally sensitive and responsible manner.

1.3 | OCD genetics

OCD is ~50% heritable (Mataix-Cols et al., 2013; Pauls, 2010; van Grootheest et al., 2005). First-degree relatives of affected individuals have a 4-8x increased risk of OCD (Browne et al., 2015; Mataix-Cols et al., 2013). OCD linkage studies (Pauls et al., 2014), and >100 candidate gene studies (Taylor, 2013), have produced inconsistent results. Regarding rare variation, two key whole exome sequencing (WES) studies have been published. First, Cappi et al. (2020) performed WES on 222 sporadic OCD cases and their unaffected parents to identify rare de novo mutations (DNMs). Results showed that DNMs predicted to damage gene function are enriched in OCD probands. Second, Halvorsen et al. (2021) reported WES results from the largest OCD cohort to date (1313 total cases, consisting of 587 trios, 41 quartets and 644 singletons). They identified an excess of inherited and de novo loss of function (LoF) variation in LoF-intolerant genes and the most significant single-gene result was *SLITRK5* ($P = 2.3 \times 10^{-6}$).

Taken together, these data support the contribution of rare coding variants to OCD genetic risk.

Regarding common variation, the first well-powered OCD GWAS from the PGC OCD Working Group was posted to medRxiv in October 2021 (Strom et al., 2021). This study was limited to EA individuals (N = 14,140 OCD cases; N = 562,117 controls), identified the first genome-wide significant locus for OCD, and strengthened previous literature suggesting a polygenic nature of this disorder. The PGC OCD Working Group is currently working on publishing a much larger OCD GWAS (~50,000 cases; ~1 M controls), but the sample remains heavily European because no large-scale collection of non-European OCD cases has been performed. We expect that LATINO and similar studies of OCD in individuals from diverse ancestries will help to accelerate gene discovery for this understudied condition.

2 | AIMS AND METHODS

LATINO's main aims are: 1. Diversify the OCD genomics sample; 2. Discover genomic loci for OCD; and 3. Fine-map OCD loci and test accuracy for PRS (Table 1). We describe in detail below the methods proposed to achieve each of these aims.

2.1 | Aim 1

2.1.1 | Cases

Cases must have an ICD-10 diagnosis of OCD (established with the MINI/MINI-KID or SCID), have at least one biological grandparent born in Latin America (Mexico, a Caribbean Island, Central America, or South America) and be between 7 and 89 years of age. A lower limit of 7 years was chosen to allow adequate phenotyping with a range of valid instruments, while an upper limit of 89 years was chosen because anything above this age is considered protected health information in the United States. Case participants can live anywhere in Latin America, the United States, or Canada, and are consented/ assessed by investigators in their respective country. All participants are included in the study regardless of psychiatric comorbidity, if they fulfill strict diagnostic criteria for OCD. All comorbidity is registered. Patients are excluded in cases of diagnostic uncertainty, such as OCD secondary to a neurological disorder or definite CNS insult, or where the differential diagnosis between OCD and an alternative condition is unclear. Cases are reviewed by site PIs, and a standing caseness panel is held weekly via videoconference to provide a forum to discuss diagnostic issues and questions.

The LATINO network has >50 case recruitment sites at clinics, universities, psychiatric hospitals, day treatment programs, and/or partial hospitalization programs in Latin America, the United States, and Canada (see Figure 1). Participants are being recruited in person at these sites or, in some cases, through a country-specific online screening process (www.latinostudy.org). Web and social media-based advertisements are facilitating recruitment of individuals with OCD who are not engaged in clinical services.

2.1.2 | Controls

Controls are inherited from other ongoing psychiatric genetics studies being conducted across Latin America, the United States, and Canada. These studies, like LATINO, are part of the NIMH-funded Ancestral Populations Network (APN). The APN comprises several projects focused on psychiatric genetics in diverse populations and provides LATINO with a resource for ancestry-matched controls screened for psychiatric disorders. The APN includes multiple relevant studies, such as the Populations Underrepresented in Mental Illness Association Studies (PUMAS) (Lopez-Jaramillo et al., 2022), the Latino Ancestry Genomic Psychiatry Cohort (LA-GPC) (Bigdeli et al., 2020; Pato et al., 2013) and the Neuropsychiatric Genetics in Mexican Populations (NeuroMex) study (Camarena et al., 2021). At least 20,000 controls with Latin American ancestry will be available for matching.

Controls are administered a screening questionnaire composed of 32 questions from well-validated interviews and screening measures for OCD, mania, psychosis, depression, anxiety disorders, and history of alcohol, nicotine and other substance use disorders (Pato et al., 2013). In addition, there is a section specific to demographic information (i.e., age, sex, and self-identified race/ethnicity), and a section specific to medical conditions and disorders, including head trauma and seizure history. Individuals reporting no lifetime symptoms indicative of OCD will be considered eligible to be matched control participants.

2.1.3 | Approvals and informed consent

Before recruiting any participants, all sites are approved to conduct this study by both the United States Department of State as well as their local Institutional Review Board (IRB). Further, all adult participants will provide informed consent in writing or by using an electronic consenting process approved by their local IRB. All minors will provide their written/ electronic consent or verbal assent (depending on their age and local IRB requirements) and written/electronic consent from their parents or legal guardians. When permitted by the local IRB, participants are compensated up to \$25 USD after completing the core assessment battery and providing a saliva sample. All participants give permission for their de-identified data to be made available to the scientific community via having the genomic and phenotype data added to repositories (e.g., the U.S. NIMH Data Archive) and shared with international consortia (e.g., PGC). When permitted by the local IRB, an aliquot of DNA will be sent to the NIMH Repository and Genomics Resource (NRGR).

2.1.4 | Saliva collection and DNA

All participants donate 2 ml of saliva for DNA. Saliva is collected into SDNA-1000 kits from Spectrum Solutions (Draper, Utah), which stabilize the DNA for shipping and storage at room temperature. Participants are instructed not to eat, drink, smoke, or chew gum for 30 min prior to collecting the sample. Participants recruited in person are provided a saliva kit for sampling during their appointment, while those recruited online are mailed a saliva kit with a prepaid return envelope.

 TABLE 2
 Harmonized core assessment measures used across all collection sites.

| Age | Diagnostic | Clinician administered | Self-report | Family-report |
|-----------|--|------------------------------------|---|---|
| Pediatric | MINI-KID, select modules of the DIAMOND | CY-BOCS-I/II, GAF, CGI-Severity | OCI-4, PROMIS Anxiety and Depression, PQ-LES-Q, PROMIS Stigma, BEAQ, CHI-T, 3-Item Loneliness Scale, ELSI | FASA, PROMIS Anxiety and Depression, Barriers to Help Seeking/Treatment |
| Adult | MINI, SCID, select modules of the DIAMOND | Y-BOCS-I/II, GAF, CGI-Severity | OCI-4, PHQ9, GAD7, Q-LES-Q-SF, Barriers to Help Seeking/Treatment, PROMIS Stigma, BEAQ, CHI-T, 3-Item Loneliness Scale, ELSI | FAS-PV |

DNA from 1 ml of saliva is extracted in an automated fashion using Maxwell RSC Stabilized Saliva DNA Kits and a Maxwell RSC 48 Instrument from Promega (Madison, Wisconsin). The remaining saliva solution is frozen at -80 degrees C.

2.1.5 | Clinical assessments

A strength of this study is the depth and consistency of phenotyping and treatment history, which will yield a genetic study powered to analyze comorbidity, symptom dimensions, and treatment response. All LATINO sites have harmonized diagnostic and outcome assessment protocols (see Table 2). This will allow, for example, genomic analyses based on particular subgroups (e.g., tic-related OCD, familial vs. sporadic, etc.) or the use of polygenic scores to predict disease severity.

All assessments were subject to a thorough translation, back translation, and cross-cultural adaptation and validation process to guarantee construct validity. All instruments were available in English. Bibliographic research was carried out to investigate the availability of each instrument in Spanish and Portuguese, and whether there were relevant reliability and validity studies.

Instruments unavailable in the languages needed for the study (Spanish or Brazilian Portuguese) were selected for a cross-cultural validation process, to guarantee the instrument's construct validity. Parameters applied were derived from the Consensus-based Standards for the Selection of Health Measurement Instruments (COSMIN) study design checklist for patient-reported outcome measurement instruments (Mokkink et al., 2019; Terwee et al., 2018). First, independent translations were performed by two investigators. Second, a consensus was made by one specialist and discordances were reviewed by a second specialist. Third, a back-translation was performed by an English-certified proficient investigator. Fourth, the back translation was evaluated by the author of the original instrument and after the author's suggestions and approval, adjustments were made to develop the final version. Fifth, the final version of the instrument was submitted to the test phase where \sim 40 participants were interviewed, even for self-report measures, about the clarity and accuracy of the instruments. Final instrument adjustments were made according to the test phase results.

The core battery includes a variety of clinician-administered, selfreport, and parent-report (for children) measures. The Mini International

Neuropsychiatric Interview (MINI) (Sheehan et al., 1998), the Structured Clinical Interview for DSM-V (SCID) (First & Williams, 2016), and MINI-KID (Sheehan et al., 2010) are focused, structured diagnostic interviews that assess DSM-5 and ICD-10 psychiatric disorders in adults and vouth. The Children's Yale-Brown Obsessive Compulsive Scale (CY-BOCS-I/II) (Scahill et al., 1997; Storch et al., 2019) and the Y-BOCS-I/II (Goodman et al., 1989; Storch et al., 2010) are clinician-rated, semistructured interviews that assess presence and severity of OCD symptoms in youth and adults. The Obsessive-Compulsive Inventory-4 (OCI-4) (Abramovitch et al., 2021) is a self-report measure that assesses obsessive-compulsive symptom dimensions in youth and adults. Modules of the Diagnostic Interview for Anxiety, Mood, and OCD and Related Neuropsychiatric Disorders (DIAMOND) (Tolin et al., 2018) are used to assess the presence of obsessive-compulsive related disorders (OCRD). The Global Assessment of Functioning (GAF) (Schwartz & Del Prete-Brown, 2003) scale is a single, clinician-rated item assessing the overall severity of psychopathology and functional impairment in the worst week of the past month. The Clinical Global Impression-Severity (CGI-S) (Guy, 1976) is a clinician-rated assessment of clinical severity.

B Neuropsychiatric WILEY 7 of 18

Self-reported depressive and anxiety symptoms are assessed with the PHQ-9 (Kroenke et al., 2001) and GAD-7 (Spitzer et al., 2006) for adults and the PROMIS Depression and Anxiety scales (youth) (Quinn et al., 2014). PROMIS Stigma (Molina et al., 2013) measures experiences with stigma related to OCD in both adults and youth. The Family Accommodation Scale-Patient Report (adults) (Wu et al., 2016) Family Accommodation Scale-Anxiety (youth) (Lebowitz and et al., 2013, 2015) are used to assess family accommodation to the OCD symptoms. The Quality of Life Enjoyment and Satisfaction Questionnaire-Short Form (Q-LES-Q-SF) (Endicott et al., 1993) and PQ-LES-Q (Endicott et al., 2006) are self-report measures that assess satisfaction across physical health, mental health, and well-being in adults and youth. The Barriers to Help Seeking/Treatment (Goodwin et al., 2002; Marques et al., 2010) examines factors that prevent one from seeking treatment. The Brief Experiential Avoidance Questionnaire (BEAQ) (Gámez et al., 2014) is included to assess experiential avoidance in both adults and youth. To measure self-reported compulsivity and associated functional impairment, the Cambridge-Chicago Compulsivity Trait Scale (CHI-T) (Chamberlain & Grant, 2018) is used. The Three-Item Loneliness Scale (Hughes et al., 2004) is a self-report

measure that captures feelings of loneliness. To address ethical, legal, and social aspects of this work, we developed a short questionnaire that measures participants' attitudes toward genetic testing for psychiatric conditions. Beyond these core assessments, a supplemental battery of measures has been collaboratively generated. These measures are administered on a rotational basis throughout the study to minimize burden while capturing important clinical constructs.

Each participant is comprehensively assessed in person or via a secure video conferencing system in the language the person feels most comfortable using. All symptom-oriented assessments are conducted by trained raters at either the recruitment site or by Baylor College of Medicine (BCM) personnel. Cross-site training includes a review of each measure and co-rating videotaped cases. For example, raters view several videotaped ratings of the (C)Y-BOCS-I/II and then independently rate several tapes in their respective language. Raters must score within 20% of the gold standard on the (C)Y-BOCS-I/II. Regular BCM-led office hours are held to complement on-site supervision. A (C)Y-BOCS-I/II expert committee comprised of a subset of site PIs was formed to review and score all (C)Y-BOCS-I/II assessment training videos to ensure uniformity in administration and scoring. These expert committee members also made themselves available for (C)Y-BOCS-I/I consultation with other PIs and assessors.

2.1.6 Demographic assessments

In addition to collecting core demographic information (e.g., age, sex, education, family composition, etc.), comprehensive data about past diagnoses and treatment history are collected. Psychotherapy history, particularly a course of exposure and response prevention, is assessed, as well as the associated treatment response. Detailed information regarding medication history is collected, including past medication trials, highest dose, duration of treatment, side effects, and degree of symptom improvement. Barriers to treatment, social impact, and perceptions of reproductive decision-making are also collected.

2.1.7 Clinical data collection and quality assurance

Phenotypic data is collected via the HIPPA-compliant Research Electronic Data Capture (REDCap), a secure web-based survey collection, direct entry platform, and database (Harris et al., 2009, 2019). Data quality is assessed by including REDCap flags and manual data checks. Automated data checks are planned using an R-Studio API with RED-Cap. Cross-site data will be compared for missing data and interquestion correlation to determine consistency. Study assessments are audio recorded and reviewed for interrater reliability. Site PIs provide (C)YBOCS-I/II and MINI-KID/MINI supervision in order to ensure accurate assessment administration and review recorded audio of (C) YBOCS-II and MINI-KID/MINI administration, as needed. When assessment and scoring remain unresolved during local site supervision, assessors will bring concerns to the expert committee to help come to a consensus about assessment scoring.

2.2 T Aim 2

2.2.1 Genotyping

At the time of this writing, our plan is to genotype all 5000 Latino OCD cases with the Illumina Global Screening Array (GSA), which has been shown to adequately capture variation >1% across all populations in the Thousand Genomes Project, and match with data from at least 10,000 Latino controls planned to be genotyped on the same array. If, however, costs substantially decrease for a more comprehensive methodology (e.g., low-pass whole genome plus exome sequencing or standard whole genome sequencing), we will seek to change our plan. This is particularly true if the controls on which we rely are interrogated with a different method. For the purposes of this study protocol, we focus on a scenario involving GSA genotyping, given that this is the current plan.

2.2.2 Admixture and tractor

"Admixed" individuals possess DNA with more than one ancestral component. Admixed individuals, including those from Latin America, have been systematically excluded from genetic studies due to the historic lack of available tools to appropriately handle their complex genomes. If not accounted for, such population substructure can compromise analyses and bias results when highly admixed samples are included alongside more homogenous individuals using standard analysis (Sul et al., 2018). In GWAS, the specific concern regarding including admixed participants is obtaining false positive associations due to alleles being at different frequencies across populations. Previous studies have attempted to control for this by using principal components (PCs) in linear or linear mixed model frameworks. However, PCs capture only broad admixture fractions, and individuals' local ancestry makeup may differ between case and control cohorts even if their global fractions are identical. Including PCs as covariates using standard GWAS methods still leaves open the possibility for false positive associations in admixed cohorts.

To conduct gene discovery in the best-calibrated manner for these Latino cohorts, we will utilize the Tractor framework (Atkinson et al., 2021). Tractor corrects for fine-scale population structure at the level of haplotypes by using local ancestry, allowing admixed samples of all mixture proportions to be readily included in large-scale collections in a well-calibrated manner. Tractor also has been shown to boost GWAS power in admixed cohorts to identify ancestry-specific signals. Furthermore, Tractor produces ancestry-specific effect size estimates and p values, which other strategies such as classic GWAS or admixture mapping cannot. These ancestry-specific effect sizes will be extremely valuable to downstream efforts, such as in fine-mapping and building genetic risk scores (see below).

2.2.3 LATINO GWAS

First, we will perform extensive quality control (QC) of the genotyping data as described in previous work from the PGC (Howard

et al., 2019; Trubetskoy et al., 2022). Second, we will match cases and controls through an iterative procedure meant to minimize the chances of population stratification. Third, we will quantify the ancestral composition of each subject to appropriately select reference panels for local ancestry inference. Fourth, we will perform imputation using reference sets from the Trans-Omics for Precision Medicine (TOPMed) program, which has whole genome sequencing data from a large, diverse set of samples. Fifth, we will identify ancestral tracts by performing local ancestry inference using RFmix_v2 (Maples et al., 2013).

These five steps will produce the input data needed to perform a case-control GWAS using Tractor. We will include standard covariates plus an estimate of global ancestry to control for any phenotypic effect due to overall admixture makeup. The results will be combined using a model that does not assume the same effect size across ancestries. Several methods have been designed for cross-ancestry meta-analysis that allow for effect size heterogeneity (e.g., MANTRA, MR-MEGA. MAMA). For chrX, we will use an additive logistic regression model with the same covariates. We will test the genome-wide distribution of the test statistic in comparison with the expected null distribution using the genomic inflation factor λ_{GC} and QQ plots. λ_{GC} quantifies the extent of the bulk inflation resulting from a combination of true polygenic signal, systematic technical bias, and population stratification (Devlin & Roeder, 1999). In order to quantify the contribution of these factors, we will use LD Score regression (Bulik-Sullivan et al., 2015), where the intercept estimates the inflation in the mean chi-square that results from confounding biases, such as cryptic relatedness or population stratification. To declare genomewide significance, we will strictly adhere to the field-standard p value threshold of 5×10^{-8} (Pe'er et al., 2008).

A major strength of this study is the depth and consistency of phenotyping across sites (Table 2), allowing us to analyze the genetics of biologically plausible subtypes (early onset, tic-related) and symptom dimensions. As opposed to the case-control GWAS, these analyses will consider both quantitative, as well as categorical data, and are restricted to cases only.

2.2.4 | Trans-ancestry GWAS with PGC OCD

Most standard GWAS meta-analysis methods assume fixed-effects models that predict variants to have the same true marginal effect size across all studies. This is not necessarily a proper assumption in transancestry meta-analyses. Although studies to date have generally shown that the causal genetic effect of a variant is constant across populations, there are multiple reasons for differences in effect size estimates across populations. For example, different LD structures across populations, differences in gene–gene interactions or geneenvironment interactions, and phenotypic differences may produce heterogeneity across cohorts. Therefore, it is advised to model this cross-cohort heterogeneity in meta-analysis by using a random effects model. This statistical approach allows for the true marginal effect size of each variant to vary across studies, testing the null hypothesis that the mean true effect size across cohorts is zero. Alternatively, a transancestral meta-analysis model that assumes variant effect sizes may differ more between more genetically diverse populations can be used. Statistical methods appropriate to test our study aims are actively in development and we will ensure that our meta-analytic approach for PGC OCD is appropriate.

LATINO will triple the current number of non-EA OCD cases available for GWAS with PGC OCD. Currently, PGC OCD has >50,000 cases, only ~2500 of which are of non-EA. We anticipate that over the next 5 years the PGC sample will grow to around 80,000 cases, with contributions from ongoing clinical collections and large biobanking efforts such as the Million Veteran Program (MVP) (Gaziano et al., 2016) and the AllofUs Research Program (All of Us Research Program Investigators et al., 2019). These latter two efforts should help to add more non-EA OCD cases to the pool available to PGC OCD, but there will still be a heavy bias toward EA individuals. Therefore, this study will help to ensure that non-EA populations are not left behind as OCD genomics advances.

2.3 | Aim 3

2.3.1 | Fine-mapping

Genome-wide significant loci are typically large (tens of kilobases to over a megabase) and include many variants with similar p values. The causal variant(s) within GWAS peaks are often difficult to separate from those that are in LD. Fine-mapping is used to decrease the search space for the causal variant(s) and facilitate follow-up or functional genomic work. Recent work (Lam et al., 2019; Ng et al., 2017; Wyss et al., 2018) has shown that genetic diversity can be used to improve fine-mapping resolution, owing to differences in the pattern of LD between causal and non-causal variants. If we assume that causal variants will have consistent effects across populations and non-causal variants will have inconsistent effects due to population-specific LD, then we would expect causal variants to be more statistically significant and less heterogeneous in a transancestry meta-analysis compared to non-causal variants (Peterson et al., 2019). In addition, haplotype block size may be smaller in the non-European ancestry groups, improving resolution.

The development of new methods for fine-mapping, particularly trans-ancestry fine-mapping, is a very active field of research at present. Different methodologies have particular strengths and weaknesses. We will utilize SuSiEx (SUm of SIngle Effects trans-ancestry), which is a new cross-population fine-mapping extension of SuSiE (Zou et al., 2022). We will fine-map any locus that reaches genomewide significance in the trans-ancestry meta-analysis or in ancestry-specific analyses. We will pull out a 100 kb contiguous region centered on the most significant SNP (50 kb on either side). This is a conservative choice, as LD generally begins to decay between SNPs separated by >25 kb (Conrad et al., 2006). We will integrate functional annotation data, since multiple studies have shown this can improve statistical fine mapping (Kichaev et al., 2014; Kichaev & Pasaniuc, 2015). There also exists substantial functional annotation data for brain tissue and brain cells (Wang et al., 2018) that we can

cal genetics B Neuropsychiat

use for this project. Many locus-specific properties will influence our ability to fine-map a particular locus at high resolution. These include the number of causal SNPs and differences between populations in minor allele frequency, LD structure and SNP density (Schaid et al., 2018).

2.3.2 | Trans-ancestry polygenic risk score analysis

Polygenic risk scores (PRS) are used to predict complex traits using genetic data. Most PRS methods are inadequate in non-EA populations because of decreased prediction accuracy, possibly due to different patterns of LD, allele frequencies, or overall genetic architecture (Martin et al., 2020). This limits their clinical use for precision medicine efforts in non-European populations. As with fine mapping, the development of new methods for trans-ancestry PRS work is a very active field of research.

A new PRS construction method, PRS-CSx, has recently been shown to improve cross-population polygenic prediction by integrating GWAS summary statistics from multiple populations. This allows for the benefits of including the largest discovery GWAS (typically on Europeans), as well as a better-matched discovery sample of more modest size. PRS-CSx (Ruan et al., 2022) couples genetic effects across populations via a shared continuous shrinkage (CS) prior, enabling more accurate effect size estimation by sharing information between summary statistics and leveraging linkage disequilibrium diversity across discovery samples. PRS-CSx has shown improved prediction performance across traits with a wide range of genetic architectures, cross-population genetic overlaps and discovery GWAS sample sizes.

Recent multi-ancestral PRS studies have demonstrated that working with limited sample sizes for non-European cohorts is still valuable, as it allows for larger than expected gains in predicting polygenic traits (Bigdeli et al., 2020; Lam et al., 2019). We anticipate that further increases in the non-European sample size will lead to ever larger gains in variance explained and, ultimately, to the clinical utility of PRS across populations.

3 | DISCUSSION

While new discoveries are taking place in the genetic understanding of OCD, most of these advances are built upon exclusively EA samples. For example, we are very encouraged by the recent discovery of the first 30 genome-wide significant hits for OCD, but this was achieved with an entirely EA sample, limiting the utility of these results in non-EA cases and exacerbating health disparities. The LATINO study provides the opportunity to meaningfully address this issue, which has critical implications for understanding of disease pathophysiology (e.g., identifying causal variants), and ensuring accurate PRS modeling. Beyond the contribution to OCD genetics research, this collaborative study may serve as a model for engaging stakeholders in research.

There is a pressing need for more accessible evidence-based treatments for OCD throughout the world, including Latin America. Therefore, one of the major goals of LATINO is capacity building. In collaboration with the International OCD Foundation, we have already established clinical consultation groups and a Spanish-language exposure and response prevention training institute. Furthermore, the Asociación Latinoamericana de Trastorno Obsesivo Compulsivo (ALTOC) and LATINO recently hosted the very first OCD conference ever held in Latin America. This event occurred in June 2022 in Cartagena, Colombia under the name "LATINO Cartagena: The 1st Latin American Congress on OCD." This meeting was a great success with >250 attendees (face-to-face and virtual), with >90% of attendees being of Latin American ancestry. One of the unique aspects of the meeting was its broad audience-a mixture of researchers, mental health professionals, students, and people with OCD and their families. Two tracks were available to participants (one for patients and families and a second for students/researchers and clinicians), which ensured that the congress had something to offer to all attendees.

We have now established this as an annual educational meeting that will rotate throughout Latin American countries. Given the significant disability associated with OCD, increasing the number of trained clinicians, and thereby improving access to evidence-based care has the potential to reduce suffering for many individuals worldwide. Furthermore, developing academic opportunities for publications, grant applications, and presentations is a core focus of this project. Mentoring relationships have and will continue to be established with experienced investigators throughout the consortium with the goal of developing the next generation of scholars and clinicians.

LATINO has several future directions. First, although our assessment battery is uniquely comprehensive in assessing obsessive-compulsive and related symptoms, a number of constructs were not included. We are hoping for the opportunity to integrate measures of cognition, anhedonia, and suicidality. Second, we hope to generate more comprehensive genomic data. We are funded to perform microarray-based genotyping but are seeking opportunities to whole exome or whole genome sequence LATINO samples. Third, we are primarily collecting samples from individual OCD cases, but are pursuing opportunities to also collect samples and data from biological parents and other family members to facilitate trio and other family-based genetic study designs. Finally, LATINO welcomes further collaboration with anyone interested in OCD in Latino individuals and is proud to provide an infrastructure to explore new research ideas to help this underserved population.

AUTHOR CONTRIBUTIONS

Data collection: all authors. Conceptualization: James J. Crowley, Eric A. Storch, Carolina Cappi, Marcos E. Ochoa-Panaifo, Renee M. Frederick, Elizabeth G. Atkinson, Manuel Mattheisen, Stacey Pereira, Karen G. Martinez-Gonzalez, Michele T. Pato, Humberto Nicolini, Hailiang Huang. Funding acquisition: James J. Crowley, Eric A. Storch, Carolina Cappi, Marcos E Ochoa-Panaifo, Elizabeth G. Atkinson, Manuel Mattheisen, Stacey Pereira, Michele T. Pato, Hailiang Huang. Data analyses: Renee M. Frederick, Jacey L. Anderberg, Stacey Pereira, Carolina Cappi. Manuscript preparation: all authors.

CROWLEY ET AL.

AFFILIATIONS

¹Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA

²Department of Psychiatry, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA

³Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, New York, USA

⁴Departamento de Psiquiatria, Universidade de São Paulo, São Paulo, São Paulo, Brazil

⁵Universidad Privada del Norte, Lima, Lima, Peru

⁶Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, Texas, USA

⁷Department of Psychology, University of South Florida, Tampa, Florida, USA

⁸Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, Texas, USA

⁹Department of Psychiatry, University of Florida College of Medicine, Gainesville, Florida, USA

¹⁰Department of Psychology and Neuroscience, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA

¹¹Hospital Psiquiátrico de Asunción, Direccion General, Asuncion, Central, Paraguay

¹²Departamento de Psiquiatría, Hospital Psiquiátrico de Asunción, Asuncion, Central, Paraguay

¹³Hospital Nina Rodrigues/Universidade Federal do Maranhão (UFMA), Sao Luis do Maranhao, Maranhao, Brazil

¹⁴Hospital Universitário Professor Edgard Santos, Serviço de Psiquiatria, Laboratório de Neuropsicofarmacologia—LANP, Universidade Federal da Bahia, Salvador, Bahia, Brazil

¹⁵Faculdade de Medicina da Bahia, Universidade Federal da Bahia, Pós-Graduação em Medicina e Saúde, Salvador, Bahia, Brazil
¹⁶Talk TOC, Ciudad de Mexico, Mexico

¹⁷The Mathison Centre for Mental Health Research & Education, Hotchkiss Brain Institute, University of Calgary, Calgary, Alberta, Canada

¹⁸Departamento de Saúde Mental da Faculdade de Medicina, Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil

¹⁹Hospital Psiquiátrico Universitario Del Valle, Cali, Valle del Cauca, Colombia

²⁰Centro TOC México, Ciudad de Mexico, Mexico

²¹Centro Interdisciplinario de Tourette, TOC, TDAH y Trastornos Asociados (CITA), Buenos Aires, Buenos Aires, Argentina

²²Department of Psychology, Rice University, Houston, Texas, USA

²³Department of Psychiatry, University of British Columbia,

Vancouver, British Columbia, Canada

²⁴Department of Psychiatry and Behavioral Sciences, SUNY Downstate Health Sciences University, Brooklyn, New York, USA

²⁵VA New York Harbor Healthcare System, Brooklyn, New York, USA
²⁶Department of Neurosciences and Mental Health, Hospital for Sick

Children, Toronto, Ontario, Canada

²⁷Department of Psychology, University of North Texas, Denton, Texas, USA ²⁸Departamento de Psiquiatria e Medicina Legal, Universidade

Federal do Paraná, Curitiba, Parana, Brazil

²⁹Department of Psychology, University of Miami, Coral Gables, Florida, USA

³⁰Instituto Nacional de Salud Mental "Honorio Delgado-Hideyo

Noguchi", Dirección de Niños y Adolescentes Lima, Lima, Peru

³¹Universidad de Valparaíso, Instituto de Fisiología Valparaiso,
 Valparaiso, Chile
 ³²Universidad Intercontinental, Ciudad de México, Ciudad de Mexico,

Mexico

³³Department of Counseling Psychology, University of Houston, Houston, Texas, USA

³⁴Department of Psychology, Florida International University, Miami, Florida, USA

³⁵Department of Psychiatry, University of Toronto, Toronto, Ontario, Canada

³⁶Instituto Nacional de Salud Mental "Honorio Delgado-Hideyo Noguchi", Oficina Ejecutiva de Investigación, Lima, Lima, Peru

³⁷School of Medicine, Universidad San Martin de Porres, Lima, Lima, Peru

³⁸Corporación Universitaria Rafael Nuñez, Cartagena, Bolivar, Colombia

³⁹Facultad de Estudios Superiores Iztacala, Tlalnepantla de Baz, Universidad Nacional Autónoma de México, Ciudad de Mexico, Mexico

⁴⁰UTOC, Ciudad de México, Ciudad de Mexico, Mexico

⁴¹Departamento de Medicina Forense e Psiquiatria, Universidade Federal do Paraná, Curitiba, Parana, Brazil

⁴²Departamento de Psiquiatria, Universidade Federal do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, Brazil

⁴³Departamento de Psiquiatria, Clínica Médica, Porto Alegre, Rio Grande do Sul, Brazil

⁴⁴Departamento de Psiquiatria, Universidade Federal de Ciências da Saúde de Porto Alegre, Porto Alegre, Rio Grande do Sul, Brazil ⁴⁵Hospital de Clínicas da Universidade Federal do Paraná, Curitiba, Parana, Brazil

⁴⁶Departamento de Psiquiatría de Adultos, Instituto Nacional de Salud Mental "Honorio Delgado-Hideyo Noguchi", Lima, Lima, Peru

⁴⁷Departamento de Psiquiatria e Medicina Legal, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Rio de Janeiro, Brazil

⁴⁸Departamento de Psiquiatria, Instituto D'Or de Pesquisa e Ensino, Rio de Janeiro, Rio de Janeiro, Brazil

⁴⁹Fundación Universitaria del Área Andina, Escuela de Salud y Ciencias del Deporte, Bogota, Bogota, Colombia

⁵⁰Department of Psychiatry, Texas Tech University Health Sciences Center El Paso, El Paso, Texas, USA

⁵¹Rogers Behavioral Health, Oconomowoc, Wisconsin, USA

⁵²SIN TOC, Guadalajara, Jalisco, Mexico

⁵³Departamento de Psiquiatría, Universidad de Puerto Rico, San Juan, Puerto Rico, USA

⁵⁴Universidad de Puerto Rico Campus de Ciências Médicas, San Juan, Puerto Rico, USA

⁵⁵Private Practice, San Salvador, El Salvador

cal genetics B Neuropsychiatric _______ WILEY_____ 11 of 18

are

by the applicable Crea

⁵⁶Department of Psychological Science, The University of Texas Rio Grande Valley, Edinburg, Texas, USA ⁵⁷Broad Institute of MIT and Harvard, Stanley Center for Psychiatric Research, Cambridge, Massachusetts, USA ⁵⁸Center for Mental Health–Cersame, School of Medicine and Health Sciences, Universidad del Rosario, Bogota, District of Colombia, Colombia ⁵⁹Instituto Nacional de Neurología y Neurocirugía Manuel Velasco Suarez, Subdirección de Psiguiatría, Ciudad de México, Ciudad de Mexico, Mexico ⁶⁰Centro de Neurociencias, Universidad de Valparaíso, Valparaiso, Chile ⁶¹Talk TOC, Ciudad de México, Ciudad de Mexico, Mexico ⁶²Baylor College of Medicine, School of Medicine, Houston, Texas, USA ⁶³Department of Psychology, Augustana College, Rock Island, Illinois, USA ⁶⁴Department of Psychology, University of Central Oklahoma, Edmond, Oklahoma, USA ⁶⁵Departamento de Investigación Clínica, Grupo Médico Carracci, Ciudad de México, Ciudad de Mexico, Mexico ⁶⁶National Institute of Mental Heatlh (NIMH). Bethesda. Maryland, USA ⁶⁷DFW OCD Treatment Specialists, Richardson, Texas, USA ⁶⁸Specialists in OCD and Anxiety Recovery (SOAR), Richardson, Texas, USA ⁶⁹Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA ⁷⁰Programa de Ansiedade, Obsessões e Compulsões, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil ⁷¹Department of Psychology, Fordham University, New York, New York, USA ⁷²School of Nursing, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA ⁷³México Sin Estres, Ciudad de México, Ciudad de Mexico, Mexico ⁷⁴Centro Médico ABC, Ciudad de México, Ciudad de Mexico, Mexico ⁷⁵Centro Interdisciplinario de Neurociencia de Valparaíso, Universidad de Valparaíso, Valparaiso, Valparaiso, Chile ⁷⁶Clínica de Salud Mental USFQ, Universidad San Francisco de Quito, Quito, Pichincha, Ecuador ⁷⁷Universidad San Francisco de Quito, Quito, Pichincha, Ecuador ⁷⁸Universidad Tegnológica Privada de Santa Cruz (UTEPSA), Santa Cruz de la Sierra, Andres Ibañez, Bolivia ⁷⁹Departamento Académico de Ciencias Dinámicas, Universidad Nacional Mayor de San Marcos, Lima, Lima, Peru ⁸⁰Departamento de Neurociências e Saúde Mental, Faculdade de Medicina da Bahia, Universidade Federal da Bahia, Salvador, Bahia, Brazil ⁸¹Faculdade Santa Casa, Faculdade de Psicologia, Salvador, Bahia, Brazil

⁸²Department of Psychiatry, Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada ⁸³Department of Psychiatry, Frederick Thompson Anxiety Disorders Centre, Toronto, Ontario, Canada

⁸⁴Department of Psychiatry, Temerty Faculty of Medicine, University of Toronto, Toronto, Ontario, Canada

⁸⁵Departamento de Psiquiatría, Universidad del Valle, Cali, Valle del Cauca, Colombia

⁸⁶Departamento de Psiquiatria, Universidad ICESI, Cali, Valle del Cauca, Colombia

⁸⁷Departamento de Psiquiatria, Fundación Valle del Lili, Cali, Valle del Cauca, Colombia

⁸⁸Departamento de Psiquiatria da, Universidade Federal de São Paulo (UNIFESP), Sao Paulo, Sao Paulo, Brazil

⁸⁹Facultad de Psicología, Universidad de Buenos Aires, Buenos Aires, Buenos Aires, Argentina

⁹⁰BC Mental Health and Substance Use Services, Vancouver, British Columbia, Canada

⁹¹BC Children's Hospital Research Institute, Vancouver, British Columbia, Canada

⁹²Washington University in St. Louis, St. Louis, Missouri, USA

⁹³Departamento de Psicología, Universidad de Monterrey, Monterrey, Nuevo Leon, Mexico

⁹⁴Hospital Nacional Arzobispo Loayza, Servicio de Salud Mental, Lima, Lima, Peru

 ⁹⁵Department of Molecular Brain Sciences, Centre for Addiction and Mental Health, Neurogenetics Section, Toronto, Ontario, Canada
 ⁹⁶Facultad de Ciencias de la Salud, Universidad ICESI, Cali, Valle, Colombia

⁹⁷Department of Neurology, Global Brain Health Institute—University of California San Francisco, San Francisco, California, USA

⁹⁸Departamento de Salud Mental, Hospital de Niños Ricardo

Gutierrez, Buenos Aires, Buenos Aires, Argentina

⁹⁹Hospital Universitario Austral, Materno Infantil, Buenos Aires, Buenos Aires, Argentina

 ¹⁰⁰Centro de Investigación en Ciencias y Humanidades, Universidad
 Dr. José Matías Delgado, Santa Tecla, La Libertad, El Salvador
 ¹⁰¹Department of Psychiatry, New York State Psychiatric Institute, New York, New York, USA

¹⁰²Department of Psychiatry and Behavioral Sciences, Stanford University School of Medicine, Stanford, California, USA

¹⁰³Department of Psychiatry, Temerty Faculty of Medicine, Veterans
 Affairs Palo Alto Health Care System, Palo Alto, California, USA
 ¹⁰⁴Centro Interdisciplinario de Neurociencia de Valparaiso (CINV),

Valparaiso, Chile

¹⁰⁵Instituto Realize, Buenos Aires, Buenos Aires, Argentina
 ¹⁰⁶Facultad de Psicología, Universidad Catolica Argentina, Buenos Aires, Buenos Aires, Argentina

¹⁰⁷Asociación de Psiquiatras Argentinos (APSA), Buenos Aires, Buenos Aires, Argentina

¹⁰⁸Asociación de Psiquiatras Argentinos (APSA), Presidente del Capítulo de Investigacion en Psiquiatria, Buenos Aires, Buenos Aires, Argentina

¹⁰⁹Department of Community Health and Epidemiology & Faculty of Computer Science, Dalhousie University, Halifax, Nova Scotia, Canada ¹¹⁰LMU Munich, Institute of Psychiatric Phenomics and Genomics (IPPG), Munich, Germany

¹¹¹Baylor College of Medicine, Center for Medical Ethics and Health Policy, Houston, Texas, USA

¹¹²Center for Bioethics, Harvard University School of Medicine, Boston, Massachusetts, USA

¹¹³Department of Psychiatry, Massachusetts General Hospital, Boston, Massachusetts, USA

¹¹⁴Department of Psychiatry, Rutgers University—Robert Wood Johnson Medical School, Piscataway, New Jersey, USA
¹¹⁵Departamento de Psiquiatría, Ciudad de México, Grupo Médico

Carracci, Ciudad de Mexico, Mexico

¹¹⁶Laboratorio de Genómica de Enfermedades Psiquiátricas y Neurodegenerativas, Ciudad de México, Instituto Nacional de Medicina Genómica, Ciudad de Mexico, Mexico

ACKNOWLEDGMENTS

The LATINO team acknowledges the contributions of Dave Keller, Esme Rosario, Natasha Sefcovic, Emily O'Bryant, Katia Regina da Silva, Cynthia Bulik, Stephanie Crowley, Leanne Scott, Brandy Duke, Katy Gathron, Duff Dean, McKenzie Sluder and Cole Whiteman for their input on various aspects of the project. We would like to acknowledge Gracielle Nonato for help translating instruments into Portuguese and members of the University of Puerto Rico Center for the Study and Treatment of Fear and Anxiety for their work on this project. Study data are collected and managed using REDCap electronic data capture tools provided by the Baylor College of Medicine (BCM). We'd also like to thank Uma Ramamurthy and her team at BCM Research IT for designing the LATINO study website. All LATINO sites thank, in particular, the essential contribution of those with lived experience and their families for participating in LATINO. CONSORTIA: Members of the Latin American Trans-ancestry INitiative for OCD genomics (LATINO): Jonathan S Abramowitz, Victor R Adorno, Cinthia Aguirre, Gilberto S Alves, Gustavo S Alves, NaEshia Ancalade, Jacey L Anderberg, Alejandro A Arellano Espinosa, Paul D Arnold, Elizabeth G Atkinson, Daphne M Ayton, Izabela G Barbosa, Laura Marcela Barón Castano, Cynthia N Barrera, María Celeste Berardo, Dayan Berrones, John R Best, Tim B Bigdeli, Tania Borda, Christie L Burton, Joseph D Buxbaum, Jennifer L Callahan, Carolina Cappi, Maria Cecília B Carneiro, Sandra L Cepeda, Evelyn Chazelle, Jessica M Chire, Macarena Churruca Munoz, Pamela Claisse Quiroz, Journa Cobite, Jonathan S Comer, Daniel L Costa, Jennifer Crosbie, James J Crowley, Victor O Cruz, Guillermo Dager, Luisa F Daza, Anabel de la Rosa-Gómez, Daniela del Río, Fernanda Z Delage, Carolina B Dreher, Lucila Fay, Tomas Fazio, Ygor A Ferrão, Gabriela M Ferreira, Edith G Figueroa, Leonardo F Fontenelle, Diego A Forero, Daniele TH Fragoso, Renee M Frederick, Bharathi S Gadad, Sheldon R Garrison, Paola Giusti-Rodriguez, Laura D Gonzalez, Andres González, Marco A González, Polaris Gonzalez-Barrios, Wayne K. Goodman, Dorothy E Grice, Jerry Guintivano, Daniel G Guttfreund, Andrew G Guzick, Matthew W Halvorsen, Joseph D Hovey, Hailiang Huang, Jonathan Irreño-Sotomonte, Reinhard Janssen-Aguilar, Matias Jensen, Alexandra Z Jimenez Reynolds, Joali Alexandra

Juárez Lujambio, Nasim Khalfe, Madison A Knutsen, Minjee Kook, Caleb Lack, Nuria Lanzagorta, Gabriel Lázaro-Muñoz, Monicke O Lima, Melanie O Longhurst, David A Lozada Martinez, Elba S Luna, Andrea H Marques, Molly S. Martinez, Mayra C Martinez Mallen, Karen G Martinez-Gonzalez, Maria de Los Angeles Matos, Manuel Mattheisen, Caitlyn E Maye, Joseph F McGuire, Gabriela Menezes, Euripedes C Miguel, Charlene Minaya, Tomás Miño, Sara M Mithani, Circe Montes de Oca, Alonso Morales-Rivero, Maria E Moreira-de-Oliveira, Olivia J Morris, Pablo R Moya, María Beatriz Moyano, Sandra I Muñoz, Zainab Naggash, Humberto Nicolini, Ambar A Núñez Bracho, Belinda E Núñez Bracho, Maria Corina Ochoa Rojas, Marcos E Ochoa-Panaifo, Luis A Olavarria Castaman, Trinidad Olivos Balmaceda, Iliana Ortega, Darpan I Patel, Michele T Pato, Ainsley K Patrick, Mariel Paz y Mino, Jose L Perales Orellana, Bárbara Perdigão Stumpf, Tamara Peregrina, Stacey Pereira, Tania Pérez Duarte, Kelly L Piacsek, Maritza Placencia, María Belén Prieto, Lucas C Quarantini, Yana Quarantini-Alvim, Iaroslava C Ramos, Renato T Ramos, Vanessa R Ramos, Kesley A Ramsey, Diana Rancourt, Elise V Ray, Margaret A Richter, Bradley C Riemann, Juan C Rivas, Carolyn I Rodriguez, Maria C Rosario, Camilo J Ruggero, Angel A Ruiz-Chow, Alejandra Ruiz-Velasco, Melisa N Sagarnaga, Aline S Sampaio, Leonardo C Saraiva, Russell J Schachar, Sophie C Schneider, Ethan J Schweissing, Laura D Seligman, Roseli G Shavitt, Keaton J Soileau, S. Evelvn Stewart, Eric A Storch, Shaina B Storch, Emily R Strouphauer, Vissente Tapia Cuevas, Kiara R Timpano, Beatriz Treviño-de la Garza, Alexie Vallejo-Silva, Javier Vargas-Medrano, María I Vásquez, Guadalupe Vidal Martinez, Saira A Weinzimmer, Andrew D Wiese, Mauricio A Yanez, Gwyneth Zai, Lina M Zapata-Restrepo, Luz M Zappa, Raguel M Zepeda-Burgos, Anthony W Zoghbi. Members of the Brazilian Obsessive-Compulsive Spectrum Disorder Working Group (GTTOC): Aline Santos Sampaio, Ana Gabriela Hounie, Ana Paula Ribeiro, Angela Rodrigues Souza e Silva, Ariane Fadel Martinho, Bárbara Luciane Perdigão Stumpf, Bianca Torres Mendonça de Melo Fadel, Camila Vale Porto, Carina Freitas Chacur, Carla Pereira Loureiro, Carolina Blaya Dreher, Carolina Cappi, Cristiane Flôres Bortoncello, Daniel Lucas da Conceição Costa, Daniele Tamae Hashimoto Fragoso, Euripedes Constantino Miguel, Fernanda Zetola Delage, Flávia Cervone, Gabriela Bezerra de Menezes, Gabriela Cirqueira de Souza Barros, Gabriela Mourão Ferreira, Gilberto Sousa Alves, Gustavo Santiago Alves, Hugo Caminha, Izabela Guimarães Barbosa, Jaqueline Cenci, Julia Fernandes Eigenheer Muhlbauer, Juliana Rigue da Silva, Leonardo Cardoso Saraiva, Leonardo Franklin da Costa Fontenelle, Lígia S. Camba Lopez, Livi Ferreira Testoni de Faro, Luan Pinheiro Domingues Moreira, Luana Dumans Laurito, Lucas de Castro Quarantini, Luciana Midori Samezima, Luiz Felipe Guimarães dos Santos Martins, Marcos Vinícius Sousa de Oliveira, Maria Alice de Mathis, Maria Carolina Paraíso Lopes, Maria Cecília Carneiro, Maria Conceição do Rosário, Maria Eduarda Ismerio Moreira de Oliveira, Marina Martorelli Pinho, Monicke de Oliveira Lima, Nádia Nagano, Patrycia Chedid Danna, Pedro Pereira Fortes da Silva, Raquel Fernandes de Jesus Cardoso, Renan Barbosa de Christo, Richard Chuquel Silveira de Avila, Rodrigo Bolsson Radins, Roseli Gedanke Shavitt, Samara dos Santos Ribeiro, Tatiane Veríssimo da

dical genetics B Neuropsychiate

Silveira Meirelles, Thays Mendes Aguiar, Theresa Kerolayne da Silva Sousa, Thiago Blanco Vieira, Vanessa Rogério Ramos, Veronica Huhne de Vasconcellos, Yana Quarantini Alvim, Ygor Arzeno Ferrão.

FUNDING INFORMATION

Data are being generated as part of the Ancestral Populations Network (APN), supported by the National Institute of Mental Health, USA: U01MH125050 (PI Crowley) and U01MH125062 (PI Storch). Dr. Zoghbi receives support from the NIMH (K23MH121669). Dr. Atkinson is supported by K01 MH121659, the Caroline Wiess Law Fund for Research in Molecular Medicine, and the ARCO Foundation Young Teacher-Investigator Fund at Baylor College of Medicine. Dr. Arnold receives funding from the Alberta Innovates Translational Health Chair in Child and Adolescent Mental Health. Dr. Comer receives research funding from NIH, NICHD, PCORI, SAMHSA, NSF, and FTX Foundation. Dr. Fontenelle is supported by the Conselho Nacional de Desenvolvimento Científico e. Tecnológico, Brazil (CNPg: Grant number 302526/2018-8); Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro, Brazil (FAPERJ; Grant number CNE E-26/203.052/2017); D'Or Institute of Research and Education, Brazil (IDOR; No grant number available); Coordenação de Aperfeicoamento de Pessoal de Nível Superior, Brazil (CAPES; No grant number available). Dr. Gonzalez-Barrios is supported by the NIH Scholar for Hispanic Clinical and Translational Research Education and Career Development (HCTRECD) program (R25MD007607). Dr. Zai is supported by the Academic Scholars Award at the Department of Psychiatry, University of Toronto. Dr. Miguel is supported by the National Institute of Developmental Psychiatry for Children and Adolescent (INPD) Grant: Fapesp 2014/50917-0.CNPg 465550/2014-2. Dr. Mattheisen is supported by U01MH125050. Dr. Rancourt receives support from the National Center for Complementary and Integrative Health (1R34AT010661-0). Dr. Grice is supported by the Beatrice and Samuel A. Seaver Foundation, the Mindworks Charitable Lead Trust, NIMH R01MH124679, and International OCD Foundation Innovator Award. No funding bodies were involved in the study design and collection, analysis or interpretation of data.

DATA AVAILABILITY STATEMENT

Our liberal data and analysis sharing principles will make phenotypic and genotype data and scripts widely available for access by other scientists to maximize the utility of our investigation. The datasets generated and analyzed will be available in the NIMH Data Archive (https:// nda.nih.gov/). Genomic data will also be available through the Psychiatric Genomics Consortium (PGC, https://pgc.unc.edu/). When permitted by the local IRB, DNA samples will be available from the NIMH Repository and Genomics Resource (https://www.nimhgenetics.org/).

FINANCIAL DISCLOSURES

Dr. Storch discloses the following relationships: consultant for Biohaven Pharmaceuticals and Brainsway; Book royalties from Elsevier, Springer, American Psychological Association, Wiley, Oxford, Kingsley, and Guilford; Stock valued at <\$5000 from NView; Research support from NIH, IOCDF, Ream Foundation, and Texas Higher Education Coordinating Board. Dr. Zoghbi is a consultant for AstraZeneca. In the past 3 years, Dr. Rodriguez has been a consultant for Biohaven Pharmaceuticals and Osmind; received research grant support from Biohaven Pharmaceuticals; and received a stipend from APA Publishing for her role as Deputy Editor at The American Journal of Psychiatry. Dr. Arnold receives research grant support from Biohaven Pharmaceuticals and is a consultant for Headversity. Dr. Comer received royalties from Macmillan Learning. Dr. Goodman consults for Biohaven Pharmaceuticals and has received royalties from NView LLC.

ORCID

James J. Crowley https://orcid.org/0000-0001-9051-1557 Diana Rancourt https://orcid.org/0000-0002-0241-1327 Paola Giusti-Rodriguez https://orcid.org/0000-0002-1921-1305 Jennifer L. Callahan https://orcid.org/0000-0001-9190-3886 Sheldon R. Garrison https://orcid.org/0000-0003-0745-629X Dorothy E. Grice https://orcid.org/0000-0002-8833-2549 Matias Jensen https://orcid.org/0000-0003-1327-031X Alonso Morales-Rivero https://orcid.org/0000-0002-9015-4390 Bárbara Perdigão Stumpf https://orcid.org/0000-0002-3512-0913 Roseli G. Shavitt https://orcid.org/0000-0001-5959-5748 Lina M. Zapata-Restrepo https://orcid.org/0000-0002-9615-4981 Stacey Pereira https://orcid.org/0000-0001-6541-3904

REFERENCES

- Abramovitch, A., Abramowitz, J. S., & McKay, D. (2021). The OCI-4: An ultra-brief screening scale for obsessive-compulsive disorder. *Journal of Anxiety Disorders*, 78, 102354. https://doi.org/10.1016/j.janxdis. 2021.102354
- Abramowitz, J. S., McKay, D., & Storch, E. A. (2017). The Wiley handbook of obsessive compulsive disorders. John Wiley & Sons. https://play. google.com/store/books/details?id=90kpDwAAQBAJ
- All of Us Research Program Investigators, Denny, J. C., Rutter, J. L., Goldstein, D. B., Philippakis, A., Smoller, J. W., Jenkins, G., & Dishman, E. (2019). The "all of us" research program. *The New England Journal of Medicine*, 381(7), 668–676. https://doi.org/10.1056/ NEJMsr1809937
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). American Psychiatric Association.
- Atkinson, E. G., Maihofer, A. X., Kanai, M., Martin, A. R., Karczewski, K. J., Santoro, M. L., Ulirsch, J. C., Kamatani, Y., Okada, Y., Finucane, H. K., Koenen, K. C., Nievergelt, C. M., Daly, M. J., & Neale, B. M. (2021). Tractor uses local ancestry to enable the inclusion of admixed individuals in GWAS and to boost power. *Nature Genetics*, *53*(2), 195–204. https://doi.org/10.1038/s41588-020-00766-y
- Benuto, L. T. (2017). Toolkit for counseling Spanish-speaking clients: Enhancing behavioral health services. Springer. https://play.google. com/store/books/details?id=QK4-DwAAQBAJ
- Bigdeli, T. B., Genovese, G., Georgakopoulos, P., Meyers, J. L., Peterson, R. E., Iyegbe, C. O., Medeiros, H., Valderrama, J., Achtyes, E. D., Kotov, R., Stahl, E. A., Abbott, C., Azevedo, M. H., Belliveau, R. A., Bevilacqua, E., Bromet, E. J., Byerley, W., Carvalho, C. B., Chapman, S. B., ... Pato, C. N. (2020). Contributions of common genetic variants to risk of schizophrenia among individuals of African and Latino ancestry. *Molecular Psychiatry*, *25*(10), 2455– 2467. https://doi.org/10.1038/s41380-019-0517-y
- Bloch, M. H., Landeros-Weisenberger, A., Rosario, M. C., Pittenger, C., & Leckman, J. F. (2008). Meta-analysis of the symptom structure of

obsessive-compulsive disorder. The American Journal of Psychiatry, 165(12), 1532–1542. https://doi.org/10.1176/appi.ajp.2008. 08020320

- Brander, G., Kuja-Halkola, R., Rosenqvist, M. A., Rück, C., Serlachius, E., Fernández de la Cruz, L., Lichtenstein, P., Crowley, J. J., Larsson, H., & Mataix-Cols, D. (2021). A population-based family clustering study of tic-related obsessive-compulsive disorder. *Molecular Psychiatry*, 26(4), 1224–1233. https://doi.org/10.1038/s41380-019-0532-z
- Browne, H. A., Hansen, S. N., Buxbaum, J. D., Gair, S. L., Nissen, J. B., Nikolajsen, K. H., Schendel, D. E., Reichenberg, A., Parner, E. T., & Grice, D. E. (2015). Familial clustering of tic disorders and obsessivecompulsive disorder. JAMA Psychiatry, 72(4), 359–366. https://doi. org/10.1001/jamapsychiatry.2014.2656
- Bulik-Sullivan, B. K., Loh, P.-R., Finucane, H. K., Ripke, S., Yang, J., Schizophrenia Working Group of the Psychiatric Genomics Consortium, Patterson, N., Daly, M. J., Price, A. L., & Neale, B. M. (2015). LD score regression distinguishes confounding from polygenicity in genomewide association studies. *Nature Genetics*, 47(3), 291–295. https://doi. org/10.1038/ng.3211
- Camarena, B., Atkinson, E. G., Baker, M., Becerra-Palars, C., Chibnik, L. B., Escamilla-Orozco, R., Jiménez-Pavón, J., Koenig, Z., Márquez-Luna, C., Martin, A. R., Morales-Cedillo, I. P., Olivares, A. M., Ortega-Ortiz, H., Rodriguez-Ramírez, A. M., Saracco-Alvarez, R., Basaldua, R. E., Sena, B. F., & Koenen, K. C. (2021). Neuropsychiatric genetics of psychosis in the Mexican population: A genome-wide association study protocol for schizophrenia, schizoaffective, and bipolar disorder patients and controls. *Complex Psychiatry*, 7(3–4), 60–70. https://doi. org/10.1159/000518926
- Cappi, C., Oliphant, M. E., Péter, Z., Zai, G., Conceição do Rosário, M., Sullivan, C. A. W., Gupta, A. R., Hoffman, E. J., Virdee, M., Olfson, E., Abdallah, S. B., Willsey, A. J., Shavitt, R. G., Miguel, E. C., Kennedy, J. L., Richter, M. A., & Fernandez, T. V. (2020). De novo damaging DNA coding mutations are associated with obsessivecompulsive disorder and overlap with Tourette's disorder and autism. *Biological Psychiatry*, 87(12), 1035–1044. https://doi.org/10.1016/j. biopsych.2019.09.029
- Chamberlain, S. R., & Grant, J. E. (2018). Initial validation of a transdiagnostic compulsivity questionnaire: The Cambridge-Chicago compulsivity trait scale. CNS Spectrums, 23(5), 340–346. https://doi.org/10.1017/ S1092852918000810
- Conrad, D. F., Jakobsson, M., Coop, G., Wen, X., Wall, J. D., Rosenberg, N. A., & Pritchard, J. K. (2006). A worldwide survey of haplotype variation and linkage disequilibrium in the human genome. *Nature Genetics*, 38(11), 1251–1260. https://doi.org/10.1038/ng1911
- Devlin, B., & Roeder, K. (1999). Genomic control for association studies. Biometrics, 55(4), 997–1004. https://doi.org/10.1111/j.0006-341x. 1999.00997.x
- Endicott, J., Nee, J., Harrison, W., & Blumenthal, R. (1993). Quality of life enjoyment and satisfaction questionnaire: A new measure. *Psychopharmacology Bulletin*, 29(2), 321–326. https://www.ncbi.nlm.nih.gov/ pubmed/8290681
- Endicott, J., Nee, J., Yang, R., & Wohlberg, C. (2006). Pediatric quality of life enjoyment and satisfaction questionnaire (PQ-LES-Q): Reliability and validity. *Journal of the American Academy of Child and Adolescent Psychiatry*, 45(4), 401–407. https://doi.org/10.1097/01.chi. 0000198590.38325.81
- Fernández de la Cruz, L., Rydell, M., Runeson, B., D'Onofrio, B. M., Brander, G., Rück, C., Lichtenstein, P., Larsson, H., & Mataix-Cols, D. (2017). Suicide in obsessive-compulsive disorder: A population-based study of 36 788 Swedish patients. *Molecular Psychiatry*, 22(11), 1626– 1632. https://doi.org/10.1038/mp.2016.115
- First, M. B., & Williams, J. B. W. (2016). SCID-5-CV: Structured clinical interview for DSM-5 disorders: Clinician version. American Psychiatric Association Publishing. http://worldcat.org/title/956516736

ical genetics B Neuropsychiatric Genetics —WILEY 15 of 18

- Fullana, M. A., Mataix-Cols, D., Caspi, A., Harrington, H., Grisham, J. R., Moffitt, T. E., & Poulton, R. (2009). Obsessions and compulsions in the community: Prevalence, interference, help-seeking, developmental stability, and co-occurring psychiatric conditions. *The American Journal* of Psychiatry, 166(3), 329–336. https://doi.org/10.1176/appi.ajp. 2008.08071006
- Gámez, W., Chmielewski, M., Kotov, R., Ruggero, C., Suzuki, N., & Watson, D. (2014). The brief experiential avoidance questionnaire: development and initial validation. *Psychological Assessment*, 26(1), 35– 45. https://doi.org/10.1037/a0034473
- Gaziano, J. M., Concato, J., Brophy, M., Fiore, L., Pyarajan, S., Breeling, J., Whitbourne, S., Deen, J., Shannon, C., Humphries, D., Guarino, P., Aslan, M., Anderson, D., LaFleur, R., Hammond, T., Schaa, K., Moser, J., Huang, G., Muralidhar, S., ... O'Leary, T. J. (2016). Million veteran program: A mega-biobank to study genetic influences on health and disease. *Journal of Clinical Epidemiology*, 70, 214–223. https://doi.org/10. 1016/j.jclinepi.2015.09.016
- Goodman, W. K., Price, L. H., Rasmussen, S. A., Mazure, C., Delgado, P., Heninger, G. R., & Charney, D. S. (1989). The Yale-Brown obsessive compulsive scale. *II. Validity. Archives of General Psychiatry*, 46(11), 1012–1016. https://doi.org/10.1001/archpsyc.1989.01810110054008
- Goodman, W. K., Storch, E. A., & Sheth, S. A. (2021). Harmonizing the neurobiology and treatment of obsessive-compulsive disorder. *The American Journal of Psychiatry*, 178(1), 17–29. https://doi.org/10.1176/appi.ajp.2020.20111601
- Goodwin, R., Koenen, K. C., Hellman, F., Guardino, M., & Struening, E. (2002). Helpseeking and access to mental health treatment for obsessive-compulsive disorder. Acta Psychiatrica Scandinavica, 106(2), 143–149. https://doi.org/10.1034/j.1600-0447.2002.01221.x
- Grassi, G., Cecchelli, C., Vignozzi, L., & Pacini, S. (2020). Investigational and experimental drugs to treat obsessive-compulsive disorder. *Journal of Experimental Pharmacology*, 12, 695–706. https://doi.org/10.2147/ JEP.S255375
- Guy, W. (1976). ECDEU assessment manual for psychopharmacology. https://play.google.com/store/books/details?id=B4mZEZQRZUEC
- Halvorsen, M., Samuels, J., Wang, Y., Greenberg, B. D., Fyer, A. J., McCracken, J. T., Geller, D. A., Knowles, J. A., Zoghbi, A. W., Pottinger, T. D., Grados, M. A., Riddle, M. A., Bienvenu, O. J., Nestadt, P. S., Krasnow, J., Goes, F. S., Maher, B., Nestadt, G., & Goldstein, D. B. (2021). Exome sequencing in obsessive-compulsive disorder reveals a burden of rare damaging coding variants. *Nature Neuroscience*, 24(8), 1071–1076. https://doi.org/10.1038/s41593-021-00876-8
- Harris, P. A., Taylor, R., Minor, B. L., Elliott, V., Fernandez, M., O'Neal, L., McLeod, L., Delacqua, G., Delacqua, F., Kirby, J., Duda, S. N., & RED-Cap Consortium. (2019). The REDCap consortium: Building an international community of software platform partners. *Journal of Biomedical Informatics*, 95, 103208. https://doi.org/10.1016/j.jbi.2019.103208
- Harris, P. A., Taylor, R., Thielke, R., Payne, J., Gonzalez, N., & Conde, J. G. (2009). Research electronic data capture (REDCap): a metadata-driven methodology and workflow process for providing translational research informatics support. *Journal of Biomedical Informatics*, 42(2), 377–381. https://doi.org/10.1016/j.jbi.2008.08.010
- Hasler, G., Pinto, A., Greenberg, B. D., Samuels, J., Fyer, A. J., Pauls, D., Knowles, J. A., McCracken, J. T., Piacentini, J., Riddle, M. A., Rauch, S. L., Rasmussen, S. A., Willour, V. L., Grados, M. A., Cullen, B., Bienvenu, O. J., Shugart, Y.-Y., Liang, K.-Y., Hoehn-Saric, R., ... OCD Collaborative Genetics Study. (2007). Familiality of factor analysisderived YBOCS dimensions in OCD-affected sibling pairs from the OCD collaborative genetics study. *Biological Psychiatry*, *61*(5), 617– 625. https://doi.org/10.1016/j.biopsych.2006.05.040
- Howard, D. M., Adams, M. J., Clarke, T.-K., Hafferty, J. D., Gibson, J., Shirali, M., Coleman, J. R. I., Hagenaars, S. P., Ward, J., Wigmore, E. M., Alloza, C., Shen, X., Barbu, M. C., Xu, E. Y., Whalley, H. C.,

16 of 18 WILEY medical genetics B Neuropsychiat

Marioni, R. E., Porteous, D. J., Davies, G., Deary, I. J., ... McIntosh, A. M. (2019). Genome-wide meta-analysis of depression identifies 102 independent variants and highlights the importance of the prefrontal brain regions. *Nature Neuroscience*, *22*(3), 343–352. https://doi.org/10.1038/s41593-018-0326-7

- Hughes, M. E., Waite, L. J., Hawkley, L. C., & Cacioppo, J. T. (2004). A short scale for measuring loneliness in large surveys: Results from two population-based studies. *Research on Aging*, 26(6), 655–672. https:// doi.org/10.1177/0164027504268574
- Karno, M., Golding, J. M., Sorenson, S. B., & Burnam, M. A. (1988). The epidemiology of obsessive-compulsive disorder in five US communities. *Archives of General Psychiatry*, 45(12), 1094–1099. https://doi.org/10. 1001/archpsyc.1988.01800360042006
- Kichaev, G., & Pasaniuc, B. (2015). Leveraging functional-annotation data in trans-ethnic fine-mapping studies. American Journal of Human Genetics, 97(2), 260–271. https://doi.org/10.1016/j.ajhg.2015.06.007
- Kichaev, G., Yang, W.-Y., Lindstrom, S., Hormozdiari, F., Eskin, E., Price, A. L., Kraft, P., & Pasaniuc, B. (2014). Integrating functional data to prioritize causal variants in statistical fine-mapping studies. *PLoS Genetics*, 10(10), e1004722. https://doi.org/10.1371/journal.pgen. 1004722
- Kroenke, K., Spitzer, R. L., & Williams, J. B. (2001). The PHQ-9: Validity of a brief depression severity measure. *Journal of General Internal Medicine*, 16(9), 606–613. https://doi.org/10.1046/j.1525-1497.2001. 016009606.x
- Lam, M., Chen, C.-Y., Li, Z., Martin, A. R., Bryois, J., Ma, X., Gaspar, H., Ikeda, M., Benyamin, B., Brown, B. C., Liu, R., Zhou, W., Guan, L., Kamatani, Y., Kim, S.-W., Kubo, M., Kusumawardhani, A. A. A. A., Liu, C.-M., Ma, H., ... Huang, H. (2019). Comparative genetic architectures of schizophrenia in east Asian and European populations. *Nature Genetics*, 51(12), 1670–1678. https://doi.org/10.1038/s41588-019-0512-x
- Lebowitz, E. R., Scharfstein, L., & Jones, J. (2015). Child-report of family accommodation in pediatric anxiety disorders: Comparison and integration with mother-report. *Child Psychiatry and Human Development*, 46(4), 501–511. https://doi.org/10.1007/s10578-014-0491-1
- Lebowitz, E. R., Woolston, J., Bar-Haim, Y., Calvocoressi, L., Dauser, C., Warnick, E., Scahill, L., Chakir, A. R., Shechner, T., Hermes, H., Vitulano, L. A., King, R. A., & Leckman, J. F. (2013). Family accommodation in pediatric anxiety disorders. *Depression and Anxiety*, 30(1), 47– 54. https://doi.org/10.1002/da.21998
- Leckman, J. F., Bloch, M. H., & King, R. A. (2009). Symptom dimensions and subtypes of obsessive-compulsive disorder: A developmental perspective. *Dialogues in Clinical Neuroscience*, 11(1), 21–33. https:// www.ncbi.nlm.nih.gov/pubmed/19432385
- Leckman, J. F., Denys, D., Simpson, H. B., Mataix-Cols, D., Hollander, E., Saxena, S., Miguel, E. C., Rauch, S. L., Goodman, W. K., Phillips, K. A., & Stein, D. J. (2010). Obsessive-compulsive disorder: A review of the diagnostic criteria and possible subtypes and dimensional specifiers for DSM-V. Depression and Anxiety, 27(6), 507–527. https://doi.org/ 10.1002/da.20669
- Lopez-Jaramillo, C., Neale, B., & Musick, A. (2022). PUMAS: Populations underrepresented in mental illness association studies project. *European Neuropsychopharmacology: The Journal of the European College of Neuropsychopharmacology*, 63, e23–e24. https://doi.org/10. 1016/j.euroneuro.2022.07.054
- Maples, B. K., Gravel, S., Kenny, E. E., & Bustamante, C. D. (2013). RFMix: A discriminative modeling approach for rapid and robust local-ancestry inference. *American Journal of Human Genetics*, 93(2), 278–288. https://doi.org/10.1016/j.ajhg.2013.06.020
- Marques, L., LeBlanc, N. J., Weingarden, H. M., Timpano, K. R., Jenike, M., & Wilhelm, S. (2010). Barriers to treatment and service utilization in an internet sample of individuals with obsessive-compulsive

symptoms. Depression and Anxiety, 27(5), 470-475. https://doi.org/ 10.1002/da.20694

- Martin, A. R., Gignoux, C. R., Walters, R. K., Wojcik, G. L., Neale, B. M., Gravel, S., Daly, M. J., Bustamante, C. D., & Kenny, E. E. (2020). Human demographic history impacts genetic risk prediction across diverse populations. *American Journal of Human Genetics*, 107(4), 788–789. https://doi.org/10.1016/j.ajhg.2020.08.020
- Martin, A. R., Kanai, M., Kamatani, Y., Okada, Y., Neale, B. M., & Daly, M. J. (2019). Clinical use of current polygenic risk scores may exacerbate health disparities. *Nature Genetics*, 51(4), 584–591. https://doi.org/10. 1038/s41588-019-0379-x
- Mataix-Cols, D., Boman, M., Monzani, B., Rück, C., Serlachius, E., Långström, N., & Lichtenstein, P. (2013). Population-based, multigenerational family clustering study of obsessive-compulsive disorder. JAMA Psychiatry, 70(7), 709–717. https://doi.org/10.1001/ jamapsychiatry.2013.3
- Mataix-Cols, D., Wooderson, S., Lawrence, N., Brammer, M. J., Speckens, A., & Phillips, M. L. (2004). Distinct neural correlates of washing, checking, and hoarding symptom dimensions in obsessivecompulsive disorder. Archives of General Psychiatry, 61(6), 564–576. https://doi.org/10.1001/archpsyc.61.6.564
- Mathers, C., & World Health Organization. (2008). The global burden of disease: 2004 update. World health Organization. https://books.google.com/books/about/The_Global_Burden_of_Disease.html?hl=& id=xrYYZ6Jcfv0C
- McGuire, J. F., Piacentini, J., Lewin, A. B., Brennan, E. A., Murphy, T. K., & Storch, E. A. (2015). A meta-analysis of cognitive behavior therapy and medication for child obsessive-compulsive disorder: Moderators of treatment efficacy, response, and remission. *Depression and Anxiety*, 32(8), 580–593. https://doi.org/10.1002/da.22389
- Mokkink, L. B., Prinsen, C. A., Patrick, D. L., Alonso, J., Bouter, L. M., de Vet, H. C. W., & Terwee, C. B. (2019). COSMIN study design checklist for patient-reported outcome measurement instruments. Amsterdam, the Netherlands, 1–32. https://gut.bmj.com/content/gutjnl/70/1/139/ DC1/embed/inline-supplementary-material-1.pdf?download=true
- Molina, Y., Choi, S. W., Cella, D., & Rao, D. (2013). The stigma scale for chronic illnesses 8-item version (SSCI-8): Development, validation and use across neurological conditions. *International Journal of Behavioral Medicine*, 20(3), 450–460. https://doi.org/10.1007/s12529-012-9243-4
- Ng, M. C. Y., Graff, M., Lu, Y., Justice, A. E., Mudgal, P., Liu, C.-T., Young, K., Yanek, L. R., Feitosa, M. F., Wojczynski, M. K., Rand, K., Brody, J. A., Cade, B. E., Dimitrov, L., Duan, Q., Guo, X., Lange, L. A., Nalls, M. A., Okut, H., ... North, K. E. (2017). Discovery and finemapping of adiposity loci using high density imputation of genomewide association studies in individuals of African ancestry: African ancestry anthropometry genetics consortium. *PLoS Genetics*, 13(4), e1006719. https://doi.org/10.1371/journal.pgen.1006719
- Norris, E. T., Wang, L., Conley, A. B., Rishishwar, L., Mariño-Ramírez, L., Valderrama-Aguirre, A., & Jordan, I. K. (2018). Genetic ancestry, admixture and health determinants in Latin America. *BMC Genomics*, 19(Suppl 8), 861. https://doi.org/10.1186/s12864-018-5195-7
- Pato, M. T., Sobell, J. L., Medeiros, H., Abbott, C., Sklar, B. M., Buckley, P. F., Bromet, E. J., Escamilla, M. A., Fanous, A. H., Lehrer, D. S., Macciardi, F., Malaspina, D., McCarroll, S. A., Marder, S. R., Moran, J., Morley, C. P., Nicolini, H., Perkins, D. O., Purcell, S. M., ... Pato, C. N. (2013). The genomic psychiatry cohort: Partners in discovery. *American Journal of Medical Genetics*. *Part B*, 162B(4), 306–312. https://doi.org/10.1002/ajmg.b.32160
- Pauls, D. L. (2010). The genetics of obsessive-compulsive disorder: A review. Dialogues in Clinical Neuroscience, 12(2), 149–163.
- Pauls, D. L., Abramovitch, A., Rauch, S. L., & Geller, D. A. (2014). Obsessive-compulsive disorder: An integrative genetic and neurobiological

perspective. Nature Reviews. Neuroscience, 15(6), 410-424. https://doi.org/10.1038/nrn3746

- Pe'er, I., Yelensky, R., Altshuler, D., & Daly, M. J. (2008). Estimation of the multiple testing burden for genomewide association studies of nearly all common variants. *Genetic Epidemiology*, 32(4), 381–385. https:// doi.org/10.1002/gepi.20303
- Perez, M. I., Limon, D. L., Candelari, A. E., Cepeda, S. L., Ramirez, A. C., Guzick, A. G., Kook, M., La Buissonniere Ariza, V., Schneider, S. C., Goodman, W. K., & Storch, E. A. (2022). Obsessive-compulsive disorder misdiagnosis among mental healthcare providers in Latin America. *Journal of Obsessive-Compulsive and Related Disorders*, 32, 100693. https://doi.org/10.1016/j.jocrd.2021.100693
- Peterson, R. E., Kuchenbaecker, K., Walters, R. K., Chen, C.-Y., Popejoy, A. B., Periyasamy, S., Lam, M., Iyegbe, C., Strawbridge, R. J., Brick, L., Carey, C. E., Martin, A. R., Meyers, J. L., Su, J., Chen, J., Edwards, A. C., Kalungi, A., Koen, N., Majara, L., ... Duncan, L. E. (2019). Genome-wide association studies in ancestrally diverse populations: Opportunities, methods, pitfalls, and recommendations. *Cell*, 179(3), 589–603. https://doi.org/10.1016/j.cell.2019.08.051
- Quinn, H., Thissen, D., Liu, Y., Magnus, B., Lai, J.-S., Amtmann, D., Varni, J. W., Gross, H. E., & DeWalt, D. A. (2014). Using item response theory to enrich and expand the PROMIS pediatric self report banks. *Health and Quality of Life Outcomes*, 12, 160. https://doi.org/10.1186/ s12955-014-0160-x
- Ruan, Y., Lin, Y.-F., Feng, Y.-C. A., Chen, C.-Y., Lam, M., Guo, Z., Stanley Global Asia Initiatives, He, L., Sawa, A., Martin, A. R., Qin, S., Huang, H., & Ge, T. (2022). Improving polygenic prediction in ancestrally diverse populations. *Nature Genetics*, 54(5), 573–580. https:// doi.org/10.1038/s41588-022-01054-7
- Ruscio, A. M., Stein, D. J., Chiu, W. T., & Kessler, R. C. (2010). The epidemiology of obsessive-compulsive disorder in the National Comorbidity Survey Replication. *Molecular Psychiatry*, 15(1), 53–63. https://doi. org/10.1038/mp.2008.94
- Sanderson, S. C., Diefenbach, M. A., Zinberg, R., Horowitz, C. R., Smirnoff, M., Zweig, M., Streicher, S., Jabs, E. W., & Richardson, L. D. (2013). Willingness to participate in genomics research and desire for personal results among underrepresented minority patients: A structured interview study. *Journal of Community Genetics*, 4(4), 469–482. https://doi.org/10.1007/s12687-013-0154-0
- Scahill, L., Riddle, M. A., McSwiggin-Hardin, M., Ort, S. I., King, R. A., Goodman, W. K., Cicchetti, D., & Leckman, J. F. (1997). Children's Yale-Brown obsessive compulsive scale: Reliability and validity. *Journal* of the American Academy of Child and Adolescent Psychiatry, 36(6), 844–852. https://doi.org/10.1097/00004583-199706000-00023
- Schaid, D. J., Chen, W., & Larson, N. B. (2018). From genome-wide associations to candidate causal variants by statistical fine-mapping. *Nature Reviews. Genetics*, 19(8), 491–504. https://doi.org/10.1038/s41576-018-0016-z
- Schwartz, R. C., & Del Prete-Brown, T. (2003). Construct validity of the global assessment of functioning scale for clients with anxiety disorder. *Psychological Reports*, 92(2), 548–550. https://doi.org/10.2466/ pr0.2003.92.2.548
- Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., Hergueta, T., Baker, R., & Dunbar, G. C. (1998). The miniinternational neuropsychiatric interview (M.I.N.I.): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *The Journal of Clinical Psychiatry*, *59*(Suppl 20), 22–33. quiz 34–57.
- Sheehan, D. V., Sheehan, K. H., Shytle, R. D., Janavs, J., Bannon, Y., Rogers, J. E., Milo, K. M., Stock, S. L., & Wilkinson, B. (2010). Reliability and validity of the Mini international neuropsychiatric interview for children and adolescents (MINI-KID). *The Journal of Clinical Psychiatry*, 71(3), 313–326. https://doi.org/10.4088/JCP.09m05305whi
- Spitzer, R. L., Kroenke, K., Williams, J. B. W., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: The GAD-7.

Archives of Internal Medicine, 166(10), 1092–1097. https://doi.org/10. 1001/archinte.166.10.1092

- Storch, E. A., McGuire, J. F., Wu, M. S., Hamblin, R., McIngvale, E., Cepeda, S. L., Schneider, S. C., Rufino, K. A., Rasmussen, S. A., Price, L. H., & Goodman, W. K. (2019). Development and psychometric evaluation of the children's Yale-brown obsessive-compulsive scale second edition. *Journal of the American Academy of Child and Adolescent Psychiatry*, 58(1), 92–98. https://doi.org/10.1016/j.jaac.2018. 05.029
- Storch, E. A., Rasmussen, S. A., Price, L. H., Larson, M. J., Murphy, T. K., & Goodman, W. K. (2010). Development and psychometric evaluation of the yale-brown obsessive-compulsive scale: Second edition. *Psychological Assessment*, 22(2), 223–232. https://doi.org/10.1037/a0018492
- Strom, N. I., Yu, D., Gerring, Z. F., Halvorsen, M. W., Abdellaoui, A., Rodriguez-Fontenla, C., Sealock, J. M., Bigdeli, T., Coleman, J. R. I., Mahjani, B., Thorp, J. G., Bey, K., Burton, C. L., Luykx, J. J., Zai, G., Askland, K. D., Barlassina, C., Nissen, J. B., Bellodi, L., ... Mattheisen, M. (2021). Genome-wide association study identifies new locus associated with OCD. *medRxiv*. https://doi.org/10.1101/2021. 10.13.21261078
- Sul, J. H., Martin, L. S., & Eskin, E. (2018). Population structure in genetic studies: Confounding factors and mixed models. *PLoS Genetics*, 14(12), e1007309. https://doi.org/10.1371/journal.pgen.1007309
- Taylor, S. (2013). Molecular genetics of obsessive-compulsive disorder: A comprehensive meta-analysis of genetic association studies. *Molecular Psychiatry*, 18(7), 799–805. https://doi.org/10.1038/mp.2012.76
- Terwee, C. B., Prinsen, C. A. C., Chiarotto, A., Westerman, M. J., Patrick, D. L., Alonso, J., Bouter, L. M., de Vet, H. C. W., & Mokkink, L. B. (2018). COSMIN methodology for evaluating the content validity of patient-reported outcome measures: A Delphi study. Quality of Life Research: An International Journal of Quality of Life Aspects of Treatment, Care and Rehabilitation, 27(5), 1159–1170. https://doi.org/10.1007/s11136-018-1829-0
- Tolin, D. F., Gilliam, C., Wootton, B. M., Bowe, W., Bragdon, L. B., Davis, E., Hannan, S. E., Steinman, S. A., Worden, B., & Hallion, L. S. (2018). Psychometric properties of a structured diagnostic interview for DSM-5 anxiety, mood, and obsessive-compulsive and related disorders. Assessment, 25(1), 3–13. https://doi.org/10.1177/ 1073191116638410
- Trubetskoy, V., Pardiñas, A. F., Qi, T., Panagiotaropoulou, G., Awasthi, S., Bigdeli, T. B., Bryois, J., Chen, C.-Y., Dennison, C. A., Hall, L. S., Lam, M., Watanabe, K., Frei, O., Ge, T., Harwood, J. C., Koopmans, F., Magnusson, S., Richards, A. L., Sidorenko, J., ... Schizophrenia Working Group of the Psychiatric Genomics Consortium. (2022). Mapping genomic loci implicates genes and synaptic biology in schizophrenia. *Nature*, 604(7906), 502–508. https://doi.org/10.1038/s41586-022-04434-5
- van Grootheest, D. S., Cath, D. C., Beekman, A. T., & Boomsma, D. I. (2005). Twin studies on obsessive-compulsive disorder: A review. Twin Research and Human Genetics: The Official Journal of the International Society for Twin Studies, 8(5), 450–458. https://doi.org/10.1375/ 183242705774310060
- Wang, D., Liu, S., Warrell, J., Won, H., Shi, X., Navarro, F. C. P., Clarke, D., Gu, M., Emani, P., Yang, Y. T., Xu, M., Gandal, M. J., Lou, S., Zhang, J., Park, J. J., Yan, C., Rhie, S. K., Manakongtreecheep, K., Zhou, H., ... Gerstein, M. B. (2018). Comprehensive functional genomic resource and integrative model for the human brain. *Science*, 362(6420), eaat8464. https://doi.org/10.1126/science.aat8464
- Wetterneck, C. T., Little, T. E., Rinehart, K. L., Cervantes, M. E., Hyde, E., & Williams, M. (2012). Latinos with obsessive-compulsive disorder: Mental healthcare utilization and inclusion in clinical trials. *Journal of Obsessive-Compulsive and Related Disorders*, 1(2), 85–97. https://doi. org/10.1016/j.jocrd.2011.12.001
- World Health Organization. (1992). The ICD-10 classification of mental and behavioural disorders: Clinical descriptions and diagnostic

B Neuropsychiatric Genetics WILEY 17 of 18 guidelines. World Health Organization. https://apps.who.int/iris/ handle/10665/37958

- Wu, M. S., Pinto, A., Horng, B., Phares, V., McGuire, J. F., Dedrick, R. F., Van Noppen, B., Calvocoressi, L., & Storch, E. A. (2016). Psychometric properties of the family accommodation scale for obsessivecompulsive disorder-patient version. *Psychological Assessment*, 28(3), 251–262. https://doi.org/10.1037/pas0000165
- Wyss, A. B., Sofer, T., Lee, M. K., Terzikhan, N., Nguyen, J. N., Lahousse, L., Latourelle, J. C., Smith, A. V., Bartz, T. M., Feitosa, M. F., Gao, W., Ahluwalia, T. S., Tang, W., Oldmeadow, C., Duan, Q., de Jong, K., Wojczynski, M. K., Wang, X.-Q., Noordam, R., ... London, S. J. (2018). Multiethnic meta-analysis identifies ancestry-specific and cross-ancestry loci for pulmonary function. *Nature Communications*, 9(1), 2976. https://doi.org/10.1038/s41467-018-05369-0
- Zou, Y., Carbonetto, P., Wang, G., & Stephens, M. (2022). Fine-mapping from summary data with the "sum of single effects" model. *PLoS Genet*ics, 18(7), e1010299. https://doi.org/10.1371/journal.pgen.1010299

How to cite this article: Crowley, J. J., Cappi, C., Ochoa-Panaifo, M. E., Frederick, R. M., Kook, M., Wiese, A. D., Rancourt, D., Atkinson, E. G., Giusti-Rodriguez, P., Anderberg, J. L., Latin American Trans-ancestry INitiative for OCD genomics, Brazilian Obsessive-Compulsive Spectrum Disorder Working Group, Abramowitz, J. S., Adorno, V. R., Aguirre, C., Alves, G. S., Alves, G. S., Ancalade, N., Arellano Espinosa, A. A., Arnold, P. D., ... Storch, E. A. (2023). Latin American Trans-ancestry INitiative for OCD genomics (LATINO): Study protocol. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, e32962. <u>https://doi.org/10.1002/</u> ajmg.b.32962